

Hydrogen sulfide biological oxidation by pure cultures of heterotrophic bacteria

Ana Novo^{1*}, Filipa Antunes², Inês Baptista², Ruben Jorge², Isabel Saraiva³, Irina Moreira¹,
Paula Lima Castro¹

¹ CBQF - Centro de Biotecnologia e Química Fina – Laboratório Associado, Escola Superior de Biotecnologia, Universidade Católica Portuguesa/Porto, Rua Dr. António Bernardino Almeida, 4200-072 Porto, Portugal.

² WeDoTech - Companhia de Ideias e Tecnologias Lda, CiDEB/ESB, Universidade Católica Portuguesa, Rua Dr. António Bernardino Almeida, 4200-072 Porto, Portugal.

³ EFACEC – Engenharia e Sistemas, S.A., R. Eng^o Frederico Ulrich - Apartado 3003 - 4471-907 Maia, Portugal

Keywords: Hydrogen sulfide; biological oxidation, *Pseudomonas* sp., sulfide oxidase

Biogas is a mixture of gases produced during anaerobic treatment of sludge whose composition depends on the type of digested materials as well as on the operating conditions of bioreactor. This consists mainly of CH₄ and CO₂, but components such as H₂S is of particular interest due to its corrosive, toxic and environmentally hazardous properties. Removal of H₂S present in biogas can be achieved through physical and chemical processes, which are effective but produce secondary waste, which in turn gives rise to another problem of pollution. Biological treatment can be used to removing H₂S using different species of microorganisms whose specific enzymes catalyze the biological oxidation of H₂S, including photoautotrophic and chemotrophs organisms.

This work aims to compare the biotechnological removal of H₂S from biogas in aerobic conditions through heterotrophic microbial biomass and biocatalytic treatments.

Heterotrophic microorganisms were isolated from microbial enrichments supplied with H₂S streams from a wastewater treatment plant and were characterized based on their ability to grow in mineral medium with acetate as source of carbon. Isolates retrieved were mainly affiliated to the class of γ -proteobacteria with a strong prevalence of the genus *Pseudomonas*. Isolates A9, B9 and C1 all identified as *Pseudomonas* spp., revealed as the species with higher potential for H₂S removal, growing at concentrations of H₂S up to 16 mM. Total protein profiles were screened by SDS-PAGE for the three isolates, before and after the addition of H₂S, with the aim of purifying enzyme fractions involved in the oxidation process.

The authors hereby acknowledge the Agency of Innovation (Agência de Inovação, ADI, Portugal) and Quadro de Referência Estratégico Nacional (QREN, Portugal) which through the project “HbioS-Inovações biotecnológicas na remoção de H₂S presente no biogás” (QREN-ADI 21592) provided funding for the realization of this work and to the project PEst-OE/EQB/LA0016/2013.

* Corresponding author: anovo@porto.ucp.pt