

## How Dietary Intake Has Been Assessed In African Countries? A Systematic Review

### Author Information

Dr Catarina Vila-Real Masters (Corresponding Author)

Email: cvreal@porto.ucp.pt

### Affiliation 1:

Universidade Católica Portuguesa, CBQF - Centro de Biotecnologia e Química Fina –  
Laboratório Associado, Escola Superior de Biotecnologia, Rua Arquiteto Lobão Vital, Apartado  
2511, 4202-401 Porto, Portugal, Porto, Portugal

-----  
Dr Ana Pimenta-Martins

Email: apimenta@porto.ucp.pt

### Affiliation 1:

Universidade Católica Portuguesa, CBQF - Centro de Biotecnologia e Química Fina –  
Laboratório Associado, Escola Superior de Biotecnologia, Rua Arquiteto Lobão Vital, Apartado  
2511, 4202-401 Porto, Portugal, Porto, Portugal

-----  
Professor Ana Maria Gomes

Email: amgomes@porto.ucp.pt

Affiliation 1:

Universidade Católica Portuguesa, CBQF - Centro de Biotecnologia e Química Fina –  
Laboratório Associado, Escola Superior de Biotecnologia, Rua Arquiteto Lobão Vital, Apartado  
2511, 4202-401 Porto, Portugal, Porto, Portugal

-----  
Professor Elisabete Pinto

Email: epinto@porto.ucp.pt

Affiliation 1:

Universidade Católica Portuguesa, CBQF - Centro de Biotecnologia e Química Fina –  
Laboratório Associado, Escola Superior de Biotecnologia, Rua Arquiteto Lobão Vital, Apartado  
2511, 4202-401 Porto, Portugal, Porto, Portugal

Affiliation 2:

Instituto de Saúde Pública, Universidade do Porto, Rua das Taipas, 135, 4050-600 Porto,  
Portugal, Porto, Portugal

-----  
Dr Ndegwa Henry Maina

Email: henry.maina@helsinki.fi

Affiliation 1:

University of Helsinki , Department of Food and Environmental Sciences, Division of Food  
Technology, Agnes Sjöbergin tie 2, P.O. Box 66, FIN-00014 Helsinki, Finland, Helsinki,  
Finland

## INTRODUCTION

Over the last years, we have witnessed a shift in demographics in developing countries, namely in what concerns the lifetime expectancy and the organization of the societies, since there has been a growth in urbanization. This reality has led to changes in people's lifestyles, resulting in a transition from traditional to modern realities, subsequently leading to an epidemiological transition. Developing countries, especially in Africa have shown an increase in the prevalence of Non-Communicable Diseases (NCDs), while Communicable Diseases are still a major challenge, despite the success of vaccination programs (Islam et al. 2014; Haregu et al. 2014; Boutayeb 2006; World Health Organization - Regional Office for Africa 2006). According to the World Health Organization (WHO) Global Status Report (2011), NCDs are responsible for almost 80% of deaths in low and middle-income countries. Although the major cause of deaths in African countries are communicable, maternal, perinatal, and nutritional diseases, NCDs are emerging in an exponential rate, being foreseen a switch of trends a switch of trends in 2030 (World Health Organization 2011).

Dietary patterns are often considered as one of the main causes of NCDs, so it is of utmost importance to describe the expectable nutritional transition, in order to quantify the impact of diet in this group of diseases. However, in developing countries this work is scarce or insufficiently documented, probably due to logistic and financial constraints. According to Pisa *et al.* (2014), another reason that justifies the scarcity of this work is the lack of reliable dietary assessment methodologies, which upholds the emergent need for the development, validation and standardization of tools for measuring and monitoring food intake in different countries (Pisa *et al.* 2014). In this regard some work has been done, namely by the Dietary Exposure (DEX) assessment group (Pisa *et al.* 2014), which addresses its research to studies on diet and cancer and other NCDs. Its main goal is to develop and to validate dietary methods to assess dietary exposures.

The assessment of dietary intake is imperative to know population's food habits, including the inadequacy prevalence of different nutrients, as well as the study of the relationships between dietary patterns and disease. Dietary assessment may be done at national, household and individual level, when approaching food supply and production, food purchases or food consumption, respectively (Gibson 2005; Thompson & Byers 1994). At the individual level, several methodologies may be used, and these may be divided into two major groups: retrospective and prospective methods. Retrospective methods comprise the twenty four hour Recall (24hR), the Food-Frequency Questionnaire (FFQ) and the Dietary History (DH), while prospective methods include Food/Weighed Records (WR) (Gibson 2005; Thompson & Byers 1994). Ideally these tools need to be reproducible and valid in order to assure the consistency and

accuracy of measurements (Willet 1998). The choice of an appropriate method will depend on the aims of the study, the population approached as well as the research team's experience.

This systematic review intends to summarize the most recent research conducted in this field in African countries, specifically in what concerns the most used methodologies and tools.

## METHODS

The literature search was conducted on MEDLINE®/PubMed in order to identify scientific papers focused on studies about dietary intake of different populations, among African countries. This research considers several African countries, from North, East, West, Central and South Africa. In order to narrow down and systematize the search in more recent literature, only original papers published between January 2005 and December 2014 were considered and specific key MeSH terms were used: *dietary intake*; *Africa*.

Several papers were identified but not all were considered for the systematic review. The inclusion criteria established were related to: the objective of the study: only studies intending to assess dietary intake; the methodology: only studies with a suitably described methodology; and the language: only papers written in English, Portuguese and Spanish. The exclusion criteria were: studies carried out among African people but not in African countries (for example, African immigrants in other countries); assessment of a particular nutrient or a specific food or food toxin; non-quantitative assessment of the diet; dietary assessment among children; and studies performed at household level. Studies that were focused on micronutrient assessment but

further evaluated the contribution of macronutrients were also considered. Figure 1 shows a flowchart representing the paper's selection procedure.

[Figure 1]

A total of 761 studies were identified in the initial search by using the combination of the key terms mentioned above. Out of these, 221 were excluded by reading the abstract while 72 were reviews that were not included in the present study. After applying exclusion criteria, 68 papers were excluded because they did not involve a quantitative analysis or they did not comply to the established objective, 127 studies were related to a single nutrient or food, 119 investigated only children, 12 were not performed at an individual level, 29 were carried out outside Africa, 14 were written in other languages than English, Portuguese or Spanish, six did not have a suitable description of the methodology or were carried out in very small and characteristic samples. By checking the reference lists of each of these papers, another six papers, which complied with the inclusion criteria, were identified. Thus, the final number of papers was 99. Tables 1, 2, 3 and 4 summarize the main methodological issues of the included studies, allowing a comprehensive comparison between them. Papers were divided in the four tables according to the method used for dietary assessment: 24hR (table 1) FFQ (table 2), both 24hR and FFQ (table 3) and WR (table 4). In each table, in addition to information about the country where the study was conducted and the year of publication, information about methodological issues, such as the study design, studied sample, dietary assessment methods and particularities are presented. Besides these aspects, the sampling methodology and main limitations of the studies were summarized in order to understand the most difficult challenges that researchers found in the

field. Furthermore, the application of innovative technologies on dietary assessment in African countries was explored.

## RESULTS

The research retrieved 99 papers (102 studies) carried out among different African populations, namely adults (men and/or women), adolescents and elderly people.

The included papers describe studies from twenty-two different countries, located in different African regions: Algeria (n=1), Egypt (n=1), Morocco (n=2), Sudan (n=1) and Tunisia (n=5) (Northern Africa), Ethiopia (n=6), Kenya (n=10), Malawi (n=1), Mozambique (n=3), Uganda (n=3), Tanzania (n=7), Zambia (n=2) and Zimbabwe (n=1) (Eastern Africa), Benin (n=4), Burkina Faso (n=4), Ghana (n=4), Mali (n=1) and Nigeria (n=2) (Western Africa), Botswana (n=2) and South Africa (n=38), (Southern Africa), Cameroon (n=3) and Democratic Republic of the Congo (n=1) (Central Africa), each region representing 10% (n=10), 32% (n=33), 15% (n=15) and 39% (n=40) and 4% (n=4) of the total sample, respectively. The huge representation of Southern Africa is caused by the high number of studies developed in South Africa, representing 37% (n=38) of the included papers. This division of African regions is based on United Nation (UN) Statistics Division.

### Dietary Assessment Methods

Almost all of these studies are cross-sectional studies, which capture the dietary practices in a specific population, at a particular point in time (Thompson & Byers 1994; Gibson 2005).

Overall, the two most used dietary assessment tools were the 24hR and the FFQ, which were applied separately or together. Among the studies, most of them (n=54) only used 24hR and in some cases the authors chose a single day of recall (n=20) (May et al. 2014; Mupere et al. 2012; Nyuar et al. 2012; Irvine et al. 2011; Huybregts et al. 2009; Ijarotimi & Keshinro 2008; Maruapula & Chapman-Novakofski 2008; Mounir et al. 2007; Charlton et al. 2005; Wiig & Smith 2007; Kamau-Mbuthia & Elmadfa 2007; Steyn & Nel 2006; Alemayehu et al. 2011; Gewa et al. 2008; Tesfaye et al. 2008; Walton et al. 2012; Steyn et al. 2011; Steyn et al. 2012; Scarcella et al. 2011; Wiesmann et al. 2009), whilst others utilized multiple recalls (n=34) (Zeba et al. 2014; Changamire et al. 2014; Oldewage-Theron et al. 2014; Koethe et al. 2013; Kolahdooz et al. 2013; Papathakis & Pearson 2012; Hansen et al. 2011; Luke et al. 2011; Becquey & Martin-Prevel 2010; Oldewage-Theron et al. 2010; Nago et al. 2010; Dapi et al. 2010; Sodjinou et al. 2009; Alaofè et al. 2009; Mostert et al. 2005; Addo et al. 2011; Kim et al. 2014; Rankin et al. 2011; Powell et al. 2013; Termote et al. 2012; Pereko et al. 2012; López et al. 2012; Heimbürger et al. 2010; Naude et al. 2011; Gibson et al. 2011; Lamri-Senhadjji et al. 2009; Sodjinou et al. 2008; O'Keefe et al. 2007; Boumaiza et al. 2012; Oldewage-Theron et al. 2008; Kennedy et al. 2009; Becquey et al. 2009; Oldewage-Theron et al. 2008; Oldewage-Theron et al. 2008), covering a range of two to eight recalls. One study did not mention the use of a 24hR, however the described procedure allows us to conclude that this was the methodology followed (Tesfaye et al. 2008). Some studies (Sodjinou et al. 2009; Luke et al. 2011; Maruapula & Chapman-Novakofski 2008; Zeba et al. 2014; Becquey & Martin-Prevel 2010; Alaofè et al. 2009; Kim et al. 2014; Alemayehu et al. 2011; Termote et al. 2012; Powell et al. 2013; Sodjinou et al. 2008; Wiesmann et al. 2009; Becquey et al. 2009) followed a validated method for collecting



interviewer-administered 24hR, the so called United States Department of Agriculture (USDA) Automated Multiple-Pass Method (AMPM), which is a computerized method that can be applied in person or by telephone. Five other studies (Oldewage-Theron et al. 2014; Namugumya & Muyanja 2011; Walton et al. 2012; Oldewage-Theron et al. 2008; Oldewage-Theron et al. 2008) mentioned other validated methods, one based on four steps developed by Gibson and Ferguson (Gibson & Ferguson 1999; Gibson 2005), and another one which is a 24hR questionnaire developed and validated by Oldewage-Theron et al. (Oldewage-theron et al. 2005). All the dietary information collected from these studies using the 24hR reference tool is summarized in table 1.

[Table 1]

In some papers (n=30), authors selected only the FFQ for the dietary assessment (Wrottesley et al. 2014; Lukmanji et al. 2013; Pretorius et al. 2012; Anderson et al. 2011; Joffe et al. 2011; Aounallah-Skhiri et al. 2011; Kruger et al. 2011; Joffe et al. 2010; Baroudi et al. 2014; Zingoni et al. 2009; Oguntibeju et al. 2007; Jackson et al. 2007; Belgnaoui & Belahsen 2006; Jordan et al. 2013; Kesa & Oldewage-Theron 2005; Sheehy et al. 2013; Merchant et al. 2005; Jackson et al. 2012; Wentzel-Viljoen et al. 2011; Botha et al. 2014; MacKeown et al. 2007; Kruger et al. 2012; Pisa et al. 2012; Joffe et al. 2012; Delport et al. 2011; Goedecke et al. 2009; Hogenkamp et al. 2008; Tessier et al. 2008; Vorster et al. 2007; Hattingh et al. 2006), using either quantitative FFQ (n=23) (Pretorius et al. 2012; Joffe et al. 2011; Joffe et al. 2010; Zingoni et al. 2009; Oguntibeju et al. 2007; Kruger et al. 2011; Wentzel-Viljoen et al. 2011; Botha et al. 2014; Kruger et al. 2012; Pisa et al. 2012; Joffe et al. 2012; Goedecke et al. 2009; Hogenkamp et al. 2008; Wrottesley et al. 2014; Anderson et al. 2011; Jackson et al. 2007; Belgnaoui &

Belahsen 2006; Kesa & Oldewage-Theron 2005; Sheehy et al. 2013; Jackson et al. 2012; Tessier et al. 2008; Vorster et al. 2007; Hattingh et al. 2006) or semi-quantitative FFQ (n=7) (Delport et al. 2011; Merchant et al. 2005; Jordan et al. 2013; Baroudi et al. 2014; Aounallah-Skhiri et al. 2011; Lukmanji et al. 2013; MacKeown et al. 2007). Several studies (Pretorius et al. 2012; Joffe et al. 2011; Joffe et al. 2010; Zingoni et al. 2009; Oguntibeju et al. 2007; Kruger et al. 2011; Wentzel-Viljoen et al. 2011; Botha et al. 2014; Kruger et al. 2012; Pisa et al. 2012; Joffe et al. 2012; Goedecke et al. 2009; Hogenkamp et al. 2008; Wrottesley et al. 2014; Vorster et al. 2007) developed in South Africa used a quantitative FFQ specific for the South African population, retrieved from two sources, namely the Transition Health and Urbanization in South Africa (THUSA) questionnaire, design by MacIntyre et al. (MacIntyre et al. 2001a; MacIntyre et al. 2001b) and the Dietary Assessment and Education Kit (DAEK) questionnaire, developed by Steyn & Senekal and launched by the Medical Research Council (MRC) (Steyn & Senekal 2005; de Villiers et al. 2006). Within the employment of quantitative food frequency questionnaires (FFQs), Jackson et al. (2007) and Anderson et al. (2011), used the questionnaire of Sharma et al. (1996), which was developed specifically for Cameroonian people, but no information about its validation was reported. Kesa & Oldewage-Theron (2005) and Hattingh et al. (2006) also mentioned that they used a previously validated questionnaire, but no reference was made to any paper where the validation study was published. Belgnaoui & Belahsen (2006) did not specify if the used FFQ was validated or not. Within semi-quantitative FFQs, a study carried out in Tunisia (Baroudi et al. 2014) used a validated FFQ developed in Italy, which was designed for a population with similar characteristics; both populations had cancer (Decarli et al. 1996; Franceschi et al. 1993). The study of MacKeown et al. (2007) used a

FFQ based on the one used in the Birth-To-Twenty study (Richter et al. 2007). Two other Tunisian studies (Aounallah-Skhiri et al. 2011; Tessier et al. 2008) used an already validated questionnaire (El Ati et al. 2004) with few modifications according to the studied population. There were other cases in which the authors developed their own quantitative questionnaires for implementation (Lukmanji et al. 2013; Jordan et al. 2013; Merchant et al. 2005; Jackson et al. 2012; Sheehy et al. 2013). For example, Sheehy et al. (2013) developed a specific FFQ for use among rural South Africans and Jackson et al. (2012) developed, validated and tested for reproducibility a FFQ for use among adults in Botswana. Jordan et al. (2013) and Merchant et al. (2005) developed semi-quantitative FFQs to assess dietary intake in Tanzanian women and in Zimbabwean population, respectively. Lukmanji et al. (2013), authors of a Tanzanian study, also developed their own semi-quantitative FFQ but gave no information about a validation study. It is possible to observe that there are few recent validated dietary assessment instruments for African populations. As mentioned, some of the studies described so far utilized questionnaires published before 2005 have been used for dietary assessment (Sharma et al. 1996; El Ati et al. 2004), in the reviewed studies. In other cases, questionnaires were obtained from the Demographic and Health Surveys in the correspondent country. Table 2 summarizes characteristics of all the studies which used an FFQ to measure the dietary intake.

[Table 2]

Several studies used a combination of both methods (n=11), (Korkalo et al. 2014; Mala et al. 2012; Namugumya & Muyanja 2011; Oldewage-Theron & Kruger 2011; Amare et al. 2012; Mbochi et al. 2012; Oldewage-theron et al. 2005; Baroudi et al. 2010; Oldewage-Theron et al. 2006; Waudou et al. 2005; Faber & Kruger 2005), as synthesized in table 3. Namugumya &

Muyanja (2011), applied 24hR aiming to study meal patterns and to assess meal quality, whereas with the application of a FFQ they intended to gather information on food selection patterns and portion sizes. Oldewage-theron et al. (2005) used the FFQ to study both quantitative and qualitative food consumption patterns and dietary intake of the respondents and they validated this FFQ using 24-hour recalls as a gold standard. This questionnaire was later used in 2006 by Oldewage-Theron et al. (2006). Korkalo et al. (2014) developed their own questionnaire and Mala et al. (2012) used a pre-existent FFQ (without mentioning its source) to gather information about frequency of food consumption while the 24hR was used to quantify the dietary intake. Amare et al. (2012), Mbochi et al. (2012), Faber & Kruger (2005) used a qualitative FFQ and 24hR to determine nutritional intake, while Oldewage-Theron & Kruger (2011) used a quantitative FFQ to assess dietary intake and food consumption patterns and a 24hR to confirm food variety and dietary intake. The questionnaire used by Amare *et al.* (2012) was based on the Hellen Keller International FFQ, previously used in Ethiopia. Baroudi et al. (2010) assessed dietary intake using a quantitative FFQ and performed 24hR in order to obtain more qualitative information, related to food brand names and food preparation methods. Waudo et al. (2005) used 24hR to assess what mothers had eaten in the preceding 24h and then applied an FFQ in order to obtain information about the types of foods commonly consumed.

[Table 3]

Besides these two retrospective methods, weighed records were also utilized, but in a much smaller number; only four studies. Hailelassie et al. (2013) and Gibson et al. (2008) used only a single day as a measuring unit while Olayiwola et al. (2012) and Abebe et al. (2008)

applied food records for three and two non-consecutive days, respectively. More detailed information about these studies is compiled in table 4.

[Table 4]

### **Methods for the Analysis of Food Intake Data**

A large range of software tools for the analysis of dietary data were mentioned in these studies. According to table 5 it can be observed that there is a preferential selection of food databases of the countries within the same African region. For instance, in Western Africa the *Software for Intake Distribution Estimation* (C-SIDE) developed by Iowa State University is commonly used, while in Eastern Africa, NutriSurvey is the mainly chosen software. In the Northern African countries, Bilnut Software was used for the majority and in the South, FoodFinder® (Grant et al. 1992) was clearly the most utilized software. Several countries had to update these tools with their own typical foods or recipes of composite dishes, when these were not available.

In Western African countries the most used nutritional programs were ESHA Food Processor® (Food Processor Diet Analysis & Fitness Software) (Addo et al. 2011; Wiig & Smith 2007; Huybregts et al. 2009; Pereko et al. 2012) and C-SIDE (Zeba et al. 2014; Becquey & Martin-Prevel 2010; Sodjinou et al. 2009; Sodjinou et al. 2008). Several authors used other softwares, such as *NutriData*, developed in California (Olayiwola et al. 2012), *Nutrition Data System for Research* (NDSR), developed by University of Minnesota (Luke et al. 2011) and *Nutrifiq*, based on the Canadian Nutrient File (Alaofè et al. 2009). A very comprehensive software system, named VBS Food Calculation System, was chosen for the Women's Dietary Diversity Project (in Burkina Faso and Mali) (Becquey et al. 2009; Kennedy et al. 2009). VBS

Food Calculation System is a set of three softwares, which include KOMEET (for food intake analysis), VBS MANAGER (nutrient composition information), ORION and FOOD GROUPS (both for nutrient intake by food group analysis).

In East Africa, almost all the studies performed therein used the *NutriSurvey Program*, which has seventeen different food databases (food composition tables from Tanzania, Kenya, Senegal, Mali and Germany among others) (Mbochi et al. 2012; Mala et al. 2012; Jordan et al. 2013; Mupere et al. 2012; Namugumya & Muyanja 2011; Kamau-Mbuthia & Elmadfa 2007; Korkalo et al. 2014). *ESHA Food Processor®* (Irvine et al. 2011; Haileslassie et al. 2013; Merchant et al. 2005), *FoodFinder®* (Steyn & Nel 2006; Steyn et al. 2011; Steyn et al. 2012), which includes the latest version of the South African Food Composition Database, NDSR (Koethe et al. 2013; Heimbürger et al. 2010), and WorldFood Dietary Assessment System (Gewa et al. 2008; Walton et al. 2012) were also used. Softwares such as Programme CANDAT (Powell et al. 2013), Food Meter UK 07 (Waudou et al. 2005) and *General Intake Estimation System*, developed by The National Food Institute, in Denmark (Hansen et al. 2011) (linked with Composition of Foods Commonly Eaten in East Africa, the UK Nutrient Databank and National Food Composition Tables and The Planning of Satisfactory Diets in Kenya), were also chosen for nutrient analysis.

Northern African countries based their nutrient analysis on *Bilnut Software* (Baroudi et al. 2010; Baroudi et al. 2014; Belgnaoui & Belahsen 2006). However, other softwares such as *ESHA Food Processor®* (Aounallah-Skhiri et al. 2011), DIAL Programme (López et al. 2012), developed by several authors from Alce Ingenieria, *Tableaux des valeurs nutritives* (Lamri-Senhadji et al. 2009), built by Souci and colleagues (2000), Dietetik®, designed for Tunisian

foods, Nutrilog®, a software with eleven different databases (Boumaiza et al. 2012) and *FoodBase Nutritional Program* (Nyuar et al. 2012) were also used in some studies.

Investigations carried out in South African countries mainly used *FoodFinder®* software (Wrottesley et al. 2014; Oldewage-Theron et al. 2014; Papathakis & Pearson 2012; Pretorius et al. 2012; Joffe et al. 2011; Oldewage-Theron et al. 2010; Zingoni et al. 2009; Oldewage-Theron et al. 2008; Oguntibeju et al. 2007; Charlton et al. 2005; Rankin et al. 2011; Oldewage-Theron & Kruger 2011; Joffe et al. 2010; Faber & Kruger 2005; Wentzel-Viljoen et al. 2011; Kruger et al. 2011; Mostert et al. 2005; O'Keefe et al. 2007; Naude et al. 2011; Kruger et al. 2012; Jackson et al. 2012; Hattingh et al. 2006; Vorster et al. 2007; Oldewage-Theron et al. 2008; Oldewage-Theron et al. 2008).

Other software databases such as NDSR (May et al. 2014), *NutriBase* (Kolahdooz et al. 2013; Sheehy et al. 2013), developed by CyberSoft (both based on USDA National Nutrient Database for Standard Reference), *Dietary Manager Program®* (Kesa & Oldewage-Theron 2005; Oldewage-Theron et al. 2006) managed by Oskar Scharf of Dietetic Services/Rand Software and Nutritionist Five (Maruapula & Chapman-Novakofski 2008) were also used.

In Central Africa, the used software tools in Cameroon were *Microdiet* (Anderson et al. 2011) and *Becel Institute Nutrition Software* (Dapi et al. 2010) and Lucille food analysis software (Termote et al. 2012) in Democratic Republic of Congo.

Some studies did not mention the use of specific software, only referring the use of food composition databases, as source of information for the nutrient analysis, whose analysis was performed with a tool, such as Microsoft® Office Excel or IBM SPSS software for example, to compute dietary data.

Generally in African countries, there is a lack of country-specific Food Composition Tables (FCTs), and the ones that have their own FCT, do not have it updated. For this reason some countries use FCTs from neighboring countries or use global databases. Examples of cited databases are: USDA Nutrient Database for Standard Reference and others developed by Food and Agriculture Organization (FAO), such as Food Composition Table for Use in Africa, West African Food Composition Table and Composition of Selected Foods from West Africa. USDA Database was the most cited database, in countries such as Burkina Faso, Cameroon, Ethiopia, Uganda, Zimbabwe, Cameroon or Botswana. All of these databases could be accessed on the International Network of Food Data Systems (INFOODS) directory.

[Table 5]

### **Sampling Methodology**

Sampling can be done using different methodologies, depending on the aim of the investigation, on the sample size, among other factors.

According to the range of studied papers, random and non-random sampling methods were used, random sampling being the most common. Some authors did not describe how the recruitment of subjects was done. Within random sampling the main methods used were stratified sampling, multi-stage sampling and simple random sampling. Non-random convenience sampling was also used.

In studies using a multi-stage sampling approach, probability proportionate to size method was frequently used in the first selection stage, i.e. in the selection of areas (rural/urban), districts, villages, communities or even quarters. Consequently households were simply randomly selected (Amare et al. 2012; Olayiwola et al. 2012; Nago et al. 2010; Sodjinou et al.



2009; Ijarotimi & Keshinro 2008; Oldewage-theron et al. 2005) or the walk method was used (Korkalo et al. 2014; Alemayehu et al. 2011), for the identification of the subject that fulfilled the inclusion criteria, within each household. Besides these walk methods, the usage of township maps to more easily select residential areas was also utilized as one of the initial methodologies of the sampling procedure (Oldewage-Theron et al. 2014; Kolahdooz et al. 2013; Oldewage-Theron & Kruger 2011; Hattingh et al. 2006). Few studies (Powell et al. 2013; Nago et al. 2010; Becquey et al. 2009) mentioned the possibility to access residential information in the studied city, which were provided by state agencies, such as the village government, Ministry of Education or the Higher Institute of Population Science. Since it was possible to differentiate segments in populations, in several cases creation of clusters (Aounallah-Skhiri et al. 2011; Kennedy et al. 2009; Jackson et al. 2012; Tessier et al. 2008; Waudo et al. 2005) or stratification of the sample (Haileslassie et al. 2013; Mbochi et al. 2012; Dapi et al. 2010; Maruapula & Chapman-Novakofski 2008; Mounir et al. 2007; Jackson et al. 2007; Steyn & Nel 2006; Kesa & Oldewage-Theron 2005; Luke et al. 2011; Anderson et al. 2011; Hogenkamp et al. 2008; Steyn et al. 2011; Powell et al. 2013; Steyn et al. 2012; Vorster et al. 2007) by environmental or individual factors, such as social strata, income, living in rural or urban area, age or sex, was performed. In some cases, participants were randomly recruited using advertisements which were placed in different and strategic locations, such as church groups, community centers and universities or even in local newspapers (Goedecke et al. 2009; O'Keefe et al. 2007; Charlton et al. 2005). Subjects were recruited in medical clinics, health centers or day care centers; in some papers random sampling was cited (Mostert et al. 2005; Lukmanji et al. 2013; Jordan et al. 2013; Oldewage-Theron et al. 2006) and in others no sampling method was mentioned (Wiig & Smith

2007; Oldewage-Theron et al. 2010; Kim et al. 2014; Heimbürger et al. 2010; Gibson et al. 2011; Oldewage-Theron et al. 2008; Belgnaoui & Belahsen 2006; Baroudi et al. 2010).

Convenience sampling was applied in several studies (Walton et al. 2012; Joffe et al. 2012; Gibson et al. 2008; Abebe et al. 2008; Termote et al. 2012). In other cases, authors only mention that the sampling method was not random (Wrottesley et al. 2014; Alaofè et al. 2009).

**DISCUSSION**

The purpose of this review was to summarize the methodologies and tools used in dietary intake assessment in African countries, in a ten year period, covering steps from the sampling to dietary data analysis.

When we seek to describe the dietary intake of a population, the first required step is to establish a representative sample. Many sampling and subject recruitment methods can be used and these were indeed reflected in the analyzed studies. The major part of the studies applied random recruitment. In the cases in which convenience sampling was performed, various segments of the population were not included and since it is not possible to calculate the total number of excluded people, it is also not possible to determine bias caused by the absence of these individuals in the sample (Gouveia de Oliveira 2009). Ideally, random methods should be used when the aim is to characterize a population. In the recruitment process, the selected approaches have to be adapted to the population socio-economical and educational capabilities in order to assure adequate response rates and to avoid constraints in participation. The authors of the studies included in this review used some strategies, such as: the description of the study objectives in the population's native language, overcoming language limitations; the possibility to give oral consent for participation in the investigation, overcoming limitations related to high rates of illiteracy; and picking enumerators or volunteers that understood very well the population and their habits aiming to reach their confidence, reducing the possibility of anxiety or suspicion that could be present in such situations (Ngo et al. 2009).

### **Dietary Assessment Methods**

Implementation of dietary assessment methods may be done in several ways, for instance face-to-face interviews, by telephone, by email, self-administrated or observation when using the weighing method. The selected implementation method is related mainly with social and economic context of the studied areas and the resources available for the research. In this review the majority of the included studies were performed with face-to-face interviews and in two cases, i.e. in South Africa and Cameroon, self-administration was used.

To understand which is the best methodology to choose according to the population and the purpose of the study, and considering that 24hR and FFQ methods rely on respondents' memory, it is important to evaluate the accuracy of memory-based reports. There is cognitive research that confirms that for general people it is easier to describe generic dietary patterns than to describe a specific dietary meal (Wirfalt 1998). According to Thompson & Byers (1994), in cross-sectional studies generally the most used methodology is the 24hR, as corroborated by this review. Furthermore, as shown in the analysis performed herein, FFQ was the second most used tool. Both have advantages and disadvantages and should be applicable in specific situations. In their review Pisa et al. (2014) also identified the same top chosen dietary assessment tools.

A single 24hR is an indepth-interview that must be administered by trained people. In some of the studies under revision the interviewer was either a nutritionist, a dietitian or a nutrition student who had been previously trained by experts to collect dietary data. Such extensive expert training of the interviewer in state-of-the-science methodology is of extreme importance for obtaining valid and reliable assessments and analyses of dietary intakes. Furthermore, the need for a broad perception and issue awareness is needed to successfully fulfill

collection of dietary data using this method which is dependent on the subject's memory. A well-trained interviewer will create the need and relaxed atmosphere for the subject, as well as ask key questions that help subjects remember their intake easily (Willet 1998). According to Wirfalt (1998), and also Smith and colleagues (Smith et al. 1991; Smith 1991), more important than closeness of time or number of assessed days are the cues presented to the respondent, which influence aspects of the memory structure that are accessed or activated. There is evidence that the presence of cues prior to method implementation and probes during the assessment, two strategies that were done in the majority of the reviewed interviews, may increase reliability given by individual dietary reports (Thompson & Byers 1994; Wirfalt 1998; Smith et al. 1991; Smith 1991). This was one of the reasons why experts in the area were selected to duly perform the analysis of dietary intake. However, not all studies across different countries could guarantee the collection of data by an expert, probably due to availability of these professionals. Another review, carried out in Africa, also pointed out this limitation (Pisa et al. 2014), referring that in some African countries there is a lack of well-trained nutritionists and dieticians, which makes it challenging, perhaps compromising, the transfer of knowledge and training of interviewers.

The Automated Multiple Pass Method (AMPM), cited by some authors as the reference tool to apply the 24hR, has been tested in different types of populations (Conway et al. 2003; Johnson et al. 1996; Conway et al. 2004; Jonnalagadda et al. 2000) and it has been used in the continuing National Health and Nutrition Examination Survey (NHANES). A single day does not represent the usual consumption because of day-to-day variation and for that reason several studies conducted multiple recalls. The more recalls are conducted, the greater similitude to usual consumption is obtained and, consequently, better accuracy is achieved. According to

Thompson & Byers (1994), the 24hR and the WR,, usually done multiple times, estimate with quantitative accuracy daily food and nutrients intake, while frequency methods, such as FFQ, are limited by their lack of quantitative accuracy. In the reviewed studies, when multiple recalls were applied, they were distributed in non-consecutive days, in order to include week-days and weekend. Ideally all the 7 days of the week should be assessed in order to better represent usual consumption and to avoid possible systematic differences on dietary intake in different days of the week (Willet 1998). However, the chosen number of days should be considered and decided considering the size of the sample, the purpose of the study, the accuracy desired, the monotony or variety of the diet, as well as the variability of nutrients and foods being assessed (Willet 1998). Yunsheng and colleagues (Ma et al. 2009) studied how many 24hR are required to describe an individual's intake and they concluded that three is the sufficient number of recalls, since with less than three significant differences in energy estimation were observed and with more than three this parameter did not significantly improve. In some cases there was no possibility of conducting a multiple recall. According to the perception of some authors, the monotony of the diet (Nyuar et al. 2012; Steyn et al. 2011) or the large number of respondents (Kamau-Mbuthia & Elmadfa 2007) meant that a single recall was enough. In other cases the lack of time and other resources, such as labour and finances were the main causes (Maruapula & Chapman-Novakofski 2008; Wiig & Smith 2007).

Validity of the 24hR is usually done by comparison between the reports of the respondents in the recall and the measures recorded or weighed by trained and expert observers. An experiment carried out in Ethiopia (Alemayehu et al. 2011) concluded that in the evaluated setting the 24hR was not an accurate substitute of WR. They concluded that the lack of

agreement regarding the number and type of foods between the two methods, caused by memory lapses, and inaccuracies in portion size estimation were the main sources of error. Nevertheless, Gewa et al. (2008) used a to a similar comparison and got different conclusions, supported by higher values of agreement coefficients. In this case, 24hR could be an acceptable alternative to weighing method, however they considered that it was necessary to improve the recall procedure.

Twenty-four hours recall does not cause a huge burden to the respondents as the food records do. Besides, the recall is less likely to modify eating behavior of the respondents, because it is implemented after they have eaten and it does not require literacy, which is necessary to perform a correct, informative and complete food record. In the studied populations this was the major strength of the 24hR and a common reason cited by authors for choosing the recalls rather than the food records. Notwithstanding, when compared with frequency methods these two methods have weaknesses in common, since they are not likely to represent the usual consumption of individuals as reflected in frequency methods.

Therefore, FFQ gives a better idea about the usual consumption because the retrospective period is larger. This period could be since the preceding seven days to the preceding year, for instance. The decision about the ideal time frame is related with two issues, the metabolism of the dietary factor being studied and the physiology/pathophysiology of the outcome (Willet 1998). If the preceding year is used as time frame the researchers assess the dietary intake throughout the whole year, covering both seasons usually referred as dry season and harvest season. When the reference time frame is shorter, the effect of seasonality is not considered,

which was a limitation mentioned by several authors. Nevertheless, some of them mentioned that seasonality probably does not induce major dietary modifications, even though seasonality is believed to have significant effects on the diet and nutritional status. In the context of Africa, especially in rural areas, seasonality is in fact an important issue since the production and consequent consumption of some foods, such as fruit, vegetables and cereals, are directly affected by weather conditions that characterize both dry and wet seasons (Savy et al. 2006; Asombang et al. 2013; Msaki & Hendriks 2014). These diet modifications can lead to different intake in some nutrients such as vitamins and fat (Mitchikpe et al. 2008; Wiesmann et al. 2009; Masibo 2013; Faber & Laubscher 2008). More than half of the studies that used FFQ as dietary assessment tool did not specify FFQ's time frame, although the preceding year was the most cited.

Some of the FFQ utilized by the reviewed authors were specifically created for those studies, hence they included the elaboration of the food list besides other steps. Regarding the food list, the way of organizing food items in a questionnaire determines the answer of the respondent. Wirfalt (1998) cited some studies that had better results regarding reproducibility and validity when food items were organized according to the type of meals they usually consumed rather than when they were organized according to food groups. Most of the reviewed studies that used FFQ organized their dietary information in terms of food groups.

As it is possible to observe from the results section, among the studies developed in other countries besides South Africa, few of them used FFQ as one of the selected tools. In these studies authors had to create a new FFQ because in countries such as Kenya, Mozambique, Uganda, Tanzania, Botswana, Morocco or Ethiopia there is no population-specific FFQ. This



shows the need for developing new food frequency tools within the majority of African countries.

Validity of a FFQ is not a very practical and easy process to perform, because it requires a noninvasive observation of total diet of the respondents during a long period, and these validation studies have not yet been done (Thompson & Byers 1994). What is currently and usually done is the comparison of results from FFQ with that from recalls and records (Thompson & Byers 1994), a process which for some authors should be called calibration instead of validation (Willet 1998).

Some of the presented studies were also tested for reproducibility. The previous referred THUSA questionnaire was tested for reproducibility by other authors (Wentzel-Viljoen et al. 2011) in a different population, with Setswana-speaking adults. They concluded that this questionnaire was reproducible.

Besides these reported cases there is still a lack of validated methods for use in a specific population, and thus the need of updating the validated dietary assessment methods across African countries is emergent. When a validation study is performed, researchers have more confidence in their method since it means that it can actually measure the aspects of diet that it was designed for (Willet 1998), as long as the study is well-performed.

### **Estimation of Food Portion Size**

Estimation of foods portion size is one of the challenging aspects of the recall tools (Willet 1998; Thompson & Byers 1994; Venter et al. 2000). In several households within rural settings it is common that all the family eat from a shared bowl, hampering the estimation process (Hudson 1995; Huybregts et al. 2008; Pisa et al. 2014). There are visual aids which are

used to help respondents to accurately report the amounts of food items consumed. In the reviewed studies several tools were used, such as household measures, food models (two-dimension or three-dimension), food photographs and pictures, containers, real food items, among others. Within the studies that used 24hR, some did not mention how this estimation was done leaving less margin to evaluate the associated effectiveness (May et al. 2014; Changamire et al. 2014; Oldewage-Theron et al. 2008; Maruapula & Chapman-Novakofski 2008; Mounir et al. 2007; Mostert et al. 2005; Pereko et al. 2012; López et al. 2012; Heimbürger et al. 2010; O’Keefe et al. 2007; Gibson et al. 2011). In fact, there is little data concerning the accuracy of portion size estimation tools. A study of Byrd-Bredbenner & Schwartz (2004) evaluated if using portion size measurement aids (PSMAs) had effect on portion size estimation accuracy, in a group of young adults. The PSMAs were two: one was a life-size card containing pictures of both tennis and golf balls and the other one were both real tennis and golf balls. They concluded that even if the estimation accuracy was improved by the use of PSMAs, estimation errors still remain. In Burkina Faso an album of food photographs was validated for use on food portion size estimation of frequently consumed food items (Huybregts et al. 2008). This validated album, with four photos per one of the eight evaluated food items, could be used in 24hR as a valuable and accurate tool in West African rural settings. Another example of advances in this area is the book of food photographs developed and tested by Venter et al. (2000), with the purpose to be used in the THUSA study. A more recent work (Lombard et al. 2013), also carried out in rural areas of South Africa, focused on the development of a food photography series, mainly geared toward oesophageal cancer patients.

In FFQs, portion size of the food items may either be or not be assessed; in this latter option it can be assumed a common portion size for all subjects. There were few papers among the many reviewed in which authors, having collected the amounts consumed by the respondents, did not report how it was done (Baroudi et al. 2014; Belgnaoui & Belahsen 2006; Botha et al. 2014; Delport et al. 2011; Tessier et al. 2008; Jackson et al. 2007; Vorster et al. 2007). The most used tools to estimate portion size were household measures, food models (two-dimension or three-dimension) and food photographs, including the validated food photo manual (Venter et al. 2000). Implementation of the FFQ by mail or by telephone was not used in the reviewed studies. Although these possibilities are considered or applied many times in European or American dietary surveys, the socio-economical, political or geophysical conditions found in many African countries may entail natural communication barriers.

### **Food Composition Databases**

To convert the dietary intake into nutrient intake some components are needed, such as a food composition database, a coding system for matching foods listed with the entries in the food composition database and a software for calculating the nutrients' composition (Willet 1998; Thompson & Byers 1994). The right choice of the nutrient database is very important because the estimation of nutrient intake is affected by it. Parameters such as the completeness regarding the included food items and evaluated nutrients are related to the constant updating of the database, so it is imperative to support the analysis on the most recent updated version available (Thompson & Byers 1994). These nutrient databases are commonly included in computer software programs that process data and calculate individual dietary intake. The choice of the software should be based on the level of specification and detail needed, on the type of foods that

are usually consumed by the studied population and on the hardware and software requirements. As mentioned above, and also noted by other reviews (Pisa et al. 2014; Ochola & Masibo 2014; Ngo et al. 2009), there are few African countries with their own FCT, and countries without their own food table need to use either FCTs from neighboring countries or FAO's FCTs, which decreases reliability of the results. This was one of the most cited limitations by the authors. In this review several softwares were mentioned by the authors, however most of them are composed by the same FCTs, which makes imperative the need of creating updated tools. An example of an effort to improve this lack of country-specific databases is the study of Becquey et al. (2009), who developed a FCT for Burkina Faso bringing together the information of three sources, namely the FCT for Mali, supplemented by the WorldFood FCT for Senegal and the USDA database. This table was complete for energy, macronutrients and eleven micronutrients. The variability within the same continent is huge, and different lifestyles and typical food patterns are found even within the same country, which makes the finding of uniformity in the FCTs quite challenging, and eventually impossible, and so the countries find themselves obliged to create their own tools. In order to fight against the current lack of updating of these tools it is necessary to join forces geared towards the development of both new and country-specific FCTs or at least to complete the existing ones.

Besides the limitations that were mentioned along this discussion, limitations related to the adopted methodologies, to self-reporting, to small size sample were also cited. Furthermore, the traditional way of cooking is another challenging question too, because household's women resort to memory and taste rather than follow standard recipes or measurements to cook their dishes, which may hamper a reliable assessment (Wojtusiak et al. 2011).

Concerning improvements in developed countries according to new-technology based dietary assessment methods, it is envisaged that, in the coming years, these innovative tools could be used in African countries. Examples of these methods are a mobile device food record (Zhu et al. 2008) and a system based on images of foods (Schap et al. 2014). Although Wojtusiak and colleagues (Wojtusiak et al. 2011) defend that some methods based on automated analysis of photos, voice recognition and use of simple graphical symbols representing food could be applied in dietary assessment in African countries, there is still a long way to go before that may become a reality. Africa is comprised by a large part of rural areas, some of which even do not have sanitation or electricity and food insecurity is one of the major problems. Africa has a particular social organization characterized by the co-existence of several ethnic groups and societies each one with its own traditions and habits, hampering its conjoint growth and balanced development.

## CONCLUSIONS AND RECOMMENDATIONS

Globally, African countries are crossing a challenging public health crisis, which coupled to both weak and poor social and governmental structure leads to major concerns related to health, food security and socio-economic issues. Aiming to counteract the double health burden, characterized by both communicable and non-communicable diseases, a major effort is emerging toward development of health policies and the planning, development and evaluation of nutritional interventional programs.

Data obtained from this review provided a better knowledge of the research works that have been developed in African countries concerning food habits of individuals, strengthening the need to apply a bigger effort in these many nations. As shown in this review, in African countries, there is a lack of periodical, accurate, reliable and country-specific methodologies to assess dietary intake in adults. Major limitations on dietary assessment in Africa were, on the one hand, the deficiency in validated and standardized methodologies to perform the dietary assessment and, on the other, the usage of country-unspecific food composition databases. So, related to the first it is necessary to proceed with validation studies and test for reliability of the used methods, in order to assure the consistency and accuracy of measurements, as well as the confidence therein. Regarding the second cited limitation there is an emergent need to improve the already existing databases by updating food data and to develop suitable country-specific ones for those countries that don't have their own food composition table.

Countries with better social, financial and health resources evidenced more activity in this field and performed more investigations, providing greater data availability. Due to distinct

social organization of the continent, with major problems, such as high rates of inadequate education, illiteracy, food insecurity and a frail global health system, the work on this field should be continued and widened to include other African countries. Once surpassed some of these basic challenges it will be desired to follow developed countries' trends in what concerns the usage of innovative tools.

## REFERENCES

Abebe, Y. et al. (2008). Inadequate intakes of dietary zinc among pregnant women from subsistence households in Sidama, Southern Ethiopia. *Public health nutrition*. **11**: 379–386.

Addo, A. et al. (2011). Food insecurity and perceived stress but not HIV infection are independently associated with lower energy intakes among lactating Ghanaian women. *Maternal Child Nutr.* **7**: 80–91.

Alaofè, H. et al. (2009). Education and improved iron intakes for treatment of mild iron-deficiency anemia in adolescent girls in southern Benin. *Food and Nutrition Bulletin*. **30**: 24–36.

Alemayehu, A., Abebe, Y. and Gibson, R.S. (2011). A 24-h recall does not provide a valid estimate of absolute nutrient intakes for rural women in southern Ethiopia. *Nutrition*. **27**: 919–924.

Amare, B. et al. (2012). Nutritional status and dietary intake of urban residents in Gondar, Northwest Ethiopia. *BMC Public Health*. **12**: 752.

Anderson, S.G. et al. (2011). Nutrient intakes and dysglycaemia in populations of West African

origin. *The British journal of nutrition*. **105**: 297–306.

Aounallah-Skhiri, H. et al. (2011). Nutrition transition among adolescents of a south-Mediterranean country: dietary patterns, association with socio-economic factors, overweight and blood pressure. A cross-sectional study in Tunisia. *Nutrition journal*. **10**: 38.

Asombang, A.W. et al. (2013). Gastric cancer in Zambian adults: A prospective case-control study that assessed dietary intake and antioxidant status by using urinary isoprostane excretion. *American Journal of Clinical Nutrition*. **97**: 1029–1035.

El Ati, J. et al. (2004). Food Frequency Questionnaire for Tunisian dietary intakes: development, reproducibility and validity. *Arab Journal for Food and Nutrition*. **5**: 10–30.

Baroudi, O. et al. (2014). Impact of lifestyle factors and nutrients intake on occurrence of gastrointestinal cancer in Tunisian population. *Tumor Biology*, **35**: 5815–5822.

Baroudi, T. et al. (2010). Dietary intakes of essential nutrients among Arab and Berber ethnic groups on rural Tunisian island. *Nutrition*. **26**: 75–81.

Becquey, E., Capon, G. and Martin-Prével, Y. (2009). Dietary Diversity as a Measure of the Micronutrient Adequacy of Women's Diets: Results from Ouagadougou Burkina Faso Site. Food And Nutrition Technic Al A Ssistance, Technical Report.

Becquey, E. and Martin-Prevel, Y. (2010). Micronutrient Adequacy of Women ' s Diet in Urban Burkina Faso Is Low. *The Journal of Nutrition*. 2079–2085.

Belgnaoui, S. and Belahsen, R. (2006). Nutrient intake and food consumption among pregnant women from an agricultural region of Morocco. *International journal of food sciences and*



*nutrition*. **57**: 19–27.

Botha, S. et al. (2014). Associations of suPAR with lifestyle and cardiometabolic risk factors. *European Journal of Clinical Investigation*. **44**: 619–626.

Boumaiza, I. et al. (2012). Relationship Between Leptin G2548A and Leptin Receptor Q223R Gene Polymorphisms and Obesity and Metabolic Syndrome Risk in Tunisian Volunteers. *Genetic Testing and Molecular Biomarkers*. **16**: 726–733.

Boutayeb, A. (2006). The double burden of communicable and non-communicable diseases in developing countries. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. **100**: 191–199.

Bunch S., Murphy, S. (1997). WorldFood Dietary Assessment System. Available at: <http://www.fao.org/infoods/infoods/software-tools/en/>.

Byrd-Bredbenner, C. and Schwartz, J. (2004). The effect of practical portion size measurement aids on the accuracy of portion size estimates made by young adults. *Journal of Human Nutrition and Dietetics*. **17**: 351–357.

Changamire, F.T. et al. (2014). Macronutrient and sociodemographic determinants of gestational weight gain among HIV-negative women in Tanzania. *Food and nutrition bulletin*. **35**: 43–50.

Charlton, K.E., Kolbe-Alexander, T.L. and Nel, J.H. (2005). Micronutrient dilution associated with added sugar intake in elderly black South African women. *European journal of clinical nutrition*, **59**: 1030–1042.

Conway, J., Ingwersen, L. and Moshfegh, A. (2003). Effectiveness of the USDA 5-step

Multiple-Pass Method to assess food intake in obese and non-obese women. *American Journal of Clinical Nutrition*. **77**: 71–78.

Conway, J.M., Ingwersen, L.A. and Moshfegh, A.J. (2004). Accuracy of Dietary Recall Using the USDA Five-Step Multiple-Pass Method in Men: An Observational Validation Study. *Journal of the American Dietetic Association*. **104**: 595–603.

CyberSoft, I. (1986). NutriBase Professional Nutrition and Fitness Software Site. Available at: <http://www.nutribase.com/>.

Dapi, L.N. et al. (2010). Energy and nutrient intakes in relation to sex and socio-economic status among school adolescents in urban Cameroon, Africa. *Public health nutrition*. **14**: 904–913.

Davison, S. and Mandible, D. (1994). The Food Processor Plus 6.02. Available at: <http://www.esha.com/products/food-processor/>.

Decarli, A. et al. (1996). Validation of a Food-Frequency Questionnaire Dietary Intakes in Cancer Studies in Italy. *AEP*. **6**: 110–118.

Delpont, R. et al. (2011). Changes in retinol-binding protein concentrations and thyroid homeostasis with nonoccupational exposure to DDT. *Environmental Health Perspectives*. **119**: 647–651.

Erhardt, J. and Gross, R. (2007). NutriSurvey. Available at: <http://www.nutrisurvey.de/>.

Faber, M. and Kruger, S. (2005). Dietary Intake , Perceptions Regarding Body Weight , and Attitudes Toward Weight Control of Normal Weight , Overweight , and Obese Black Females. *Ethnicity*. **15**: 238-245.

Faber, M. and Laubscher, R. (2008). Seasonal availability and dietary intake of beta-carotene-rich vegetables and fruit of 2-year-old to 5-year-old children in a rural South African setting growing these crops at household level. *International journal of food sciences and nutrition*. **59**: 46–60.

Fletcher, L. (1994). Microdiet. Available at: <http://www.microdiet.co.uk/>.

Franceschi, S. et al. (1993). Reproducibility of an Italian food frequency questionnaire for cancer studies: results for specific food items. *European journal of cancer*. **29A**: 2298–2305.

Gewa, C., Murphy, S.P. and Neumann, C.G. (2008). A comparison of weighed and recalled intakes for schoolchildren and mothers in rural Kenya. *Public health nutrition*. **12**: 1197–1204.

Gibson, R.S. et al. (2011). Plasma selenium concentrations in pregnant women in two countries with contrasting soil selenium levels. *Journal of Trace Elements in Medicine and Biology*. **25**: 230–235.

Gibson, R.S. (2005). Principles of Nutrition Assessment. Second Edition, Oxford University Press, New York.

Gibson, R.S. et al. (2008). Zinc, Gravida, Infection, and Iron, but not Vitamin B-12 or Folate Status, Predict Hemoglobin during Pregnancy in Southern Ethiopia. *J Nutr.* **138**: 581–586.

Gibson, R.S. and Ferguson, E.L. (1999). An interactive 24-Hour recall for assessing the adequacy of iron and zinc intakes in developing countries. International Life Science Institute Press, Washington DC.

Godin, G. (2007). CANDAT. Available at: <http://www.foodresearch.ca/>.

Goedecke, J.H. et al. (2009). Differential effects of abdominal adipose tissue distribution on insulin sensitivity in black and white South African women. *Obesity (Silver Spring, Md.)*. **17**: 1506–1512.

Gouveia de Oliveira, A. (2009). Bioestatística, Epidemiologia e Investigação: Teoria e Aplicações Lidel, Lisboa.

Grant, K., Langenhoven, M. and Stockton, M. (1992). Foodfinder dietary analysis software. Available at: <http://safoods.mrc.ac.za/FoodfinderDEMO.pdf>

Hailelassie, K., Mulugeta, A. and Girma, M. (2013). Feeding practices, nutritional status and associated factors of lactating women in Samre Woreda, South Eastern Zone of Tigray, Ethiopia. *Nutrition journal*. **12**: 28.

Hansen, A.W. et al. (2011). Dietary patterns, food and macronutrient intakes among adults in three ethnic groups in rural Kenya. *Public health nutrition*. **14**: 1671–1679.

Haregu, T.N. et al. (2014). National Responses to HIV / AIDS and Non-Communicable Diseases in Developing Countries : Analysis of Strategic Parallels and Differences. *Journal of Public Health Research*. **3**: 25–42.

Hattingh, Z. et al. (2006). Macronutrient intake of HIV-seropositive women in Mangaung, South Africa. *Nutrition Research*. **26**: 53–58.

Heimbürger, D.C. et al. (2010). Serum phosphate predicts early mortality in adults starting antiretroviral therapy in Lusaka, Zambia: A prospective cohort study. *PLoS ONE*. **5**:1-7.

Hogenkamp, P.S. et al. (2008). Association between consumption of black tea and iron status in

adult Africans in the North West Province: the THUSA study. *The British journal of nutrition*. **100**: 430–437.

Hudson, G.J. (1995). Food intake in a west African village. Estimation of food intake from a shared bowl. *The British journal of nutrition*. **73**: 551–569.

Huybregts, L. et al. (2008). Validity of photographs for food portion estimation in a rural West African setting. *Public health nutrition*. **11**: 581–587.

Huybregts, L.F. et al. (2009). Dietary behaviour, food and nutrient intake of pregnant women in a rural community in Burkina Faso. *Maternal & child nutrition*. **5**: 211–222.

Ijarotimi, O.S. and Keshinro, O.O. (2008). Nutritional knowledge, nutrients intake and nutritional status of hypertensive patients in Ondo State, Nigeria. *Tanzania journal of health research*. **10**: 59–67.

Iowa State University. (1996). A user's guide to C-Side: software for intake distribution estimation. Available at: <http://www.card.iastate.edu/publications/synopsis.aspx?id=168>

Irvine, S.L., Hummelen, R. and Hekmat, S. (2011). Probiotic yogurt consumption may improve gastrointestinal symptoms, productivity, and nutritional intake of people living with human immunodeficiency virus in Mwanza, Tanzania. *Nutrition Research*. **31**: 875–881.

Islam, S. et al. (2014). Non Communicable Diseases (NCDs) in developing countries: a symposium report. *Globalization and Health*, **10**.

Jackson, M. et al. (2007). Diet and overweight and obesity in populations of African origin: Cameroon, Jamaica and the UK. *Public health nutrition*. **10**: 122–130.

Jackson, M.D. et al. (2012). Validation and reproducibility of an FFQ for use among adults in Botswana. *Public health nutrition*. **16**: 1995–2004.

Joffe, Y.T. et al. (2011). The -308 G/A polymorphism of the tumour necrosis factor- $\alpha$  gene modifies the association between saturated fat intake and serum total cholesterol levels in white South African women. *Genes and Nutrition*. **6**: 353–359.

Joffe, Y.T. et al. (2012). The tumor necrosis factor- $\alpha$  gene -238 G>A polymorphism, dietary fat intake, obesity risk and serum lipid concentrations in black and white South African women. *European Journal of Clinical Nutrition*. **66**: 1295–1302.

Joffe, Y.T. et al. (2010). Tumor Necrosis Factor- $\alpha$  Gene -308 G / A Polymorphism Modulates the Relationship between Dietary Fat Intake, Serum Lipids, and Obesity Risk in Black South African Women 1, 2. *The Journal of Nutrition*. 901–907.

Johnson, R.K., Driscoll, P. and Goran, M.I. (1996). Comparison of multiple-pass 24-hour recall estimates of energy intake with total energy expenditure determined by the doubly labeled water method in young children. *Journal of the American Dietetic Association*. **96**: 1140–1144.

Jonnalagadda, S.S. et al. (2000). Accuracy of energy intake data estimated by a multiple-pass, 24-hour dietary recall technique. *Journal of the American Dietetic Association*. **100**: 303–311.

Jordan, I. et al. (2013). Dietary patterns and breast cancer risk among women in northern Tanzania: A case-control study. *European Journal of Nutrition*. **52**: 905–915.

Kamau-Mbuthia, E. and Elmadfa, I. (2007). Diet quality of pregnant women attending an antenatal clinic in Nakuru, Kenya. *Annals of Nutrition and Metabolism*. **51**: 324–330.

Kennedy, G., Fanou, N. and Brouwer, I.D. (2009). Dietary Diversity as a Measure of the Micronutrient Adequacy of Women's Diets: Results from Bamako, Mali Site. Food And Nutrition Technic Al A Ssistance, Technical Report.

Kesa, H. and Oldewage-Theron, W. (2005). Anthropometric indications and nutritional intake of women in the Vaal Triangle, South Africa. *Public Health*. **119**: 294–300.

Kim, F. et al. (2014). Deficiencies of Macronutrient Intake Among HIV-Positive Breastfeeding Women in Dar es Salaam , Tanzania. **67**: 569–572.

Koethe, J.R. et al. (2013). Self-reported Dietary Intake and Appetite Predict Early Treatment Outcome among Low Body Mass Index Adults Initiating HIV Treatment in sub-Saharan Africa. *Public Helath Nutr..* **16**: 549–558.

Kolahdooz, F., Spearing, K. & Sharma, S. (2013). Dietary Adequacies among South African Adults in Rural KwaZulu-Natal. *PLoS ONE*. **8**: 1-6.

Korkalo, L. et al. (2014). A cross-sectional study on the diet and nutritional status of adolescent girls in zambia province, mozambique (the zane study): Design, methods, and population characteristics. *Journal of Medical Internet Research*. **16**: 1–13.

Kruger, A. et al. (2012). Sex differences independent of other psycho-sociodemographic factors as a predictor of body mass index in black South African adults. *Journal of Health, Population and Nutrition*. **30**: 56–65.

Kruger, A., Lekalakalamokgela, S.E. and Wentzel-Viljoen, E. (2011). Rural and Urban Older African Caregivers Coping With HIV/AIDS are Nutritionally Compromised. *Journal of nutrition*

*in gerontology and geriatrics*. **30**: 274–290.

Lamri-Senhadj, M.Y. et al. (2009). Assessment of dietary consumption and time-course of changes in serum lipids and lipoproteins before, during and after Ramadan in young Algerian adults. *Singapore Medical Journal*. **50**: 288–294.

Lombard, M. et al. (2013). A Food Photograph Series for Identifying Portion Sizes of Culturally Specific Dishes in Rural Areas with High Incidence of Oesophageal Cancer. *Nutrients*. **5**: 3118–3130.

López, P.M. et al. (2012). Nutritional Status of Adolescents in the Context of the Moroccan Nutritional Transition: the Role of Parental Education. *Journal of Biosocial Science*, **44**: 481–494.

Luke, A. et al. (2011). Protocol for the modeling the epidemiologic transition study: a longitudinal observational study of energy balance and change in body weight, diabetes and cardiovascular disease risk. *BMC Public Health*, **11**: 927.

Lukmanji, Z. et al. (2013). Dietary patterns, nutrient intake, and sociodemographic characteristics in HIV-infected Tanzanian pregnant women. *Ecology of food and nutrition*. **52**: 34–62.

Ma, Y. et al. (2009). Number of 24-Hour Diet Recalls Needed to Estimate Energy Intake. *Ann Epidemiol.*, **19**: 553–559.

MacIntyre, U.E., Venter, C.S. and Vorster, H.H. (2001)a. A culture-sensitive quantitative food frequency questionnaire used in an African population: 1. Development and reproducibility.



*Public health nutrition*. **4**: 53–62.

MacIntyre, U.E., Venter, C.S. and Vorster, H.H. (2001)b. A culture-sensitive quantitative food frequency questionnaire used in an African population: 2. Relative validation by 7-day weighted records and biomarkers. *Public health nutrition*. **4**: 63–71.

MacKeown, J.M., Pedro, T.M. and Norris, S. (2007). Energy, macro- and micronutrient intake among a true longitudinal group of South African adolescents at two interceptions (2000 and 2003): the Birth-to-Twenty (Bt20) Study. *Public health nutrition*, **10**: 635–643.

Mala, J., Tuitoek P.J. and Odhiambo R.A. (2012). Effect of Dietary Intakes on Pregnancy Outcomes: A Comparative Study among HIV-Infected and Uninfected Women at Nyanza Provincial General Hospital, Kenya. *African Journal of Food, Agriculture, Nutrition and Development*. **12**: 6776-6793.

Maruapula, S.D. and Chapman-Novakofski, K.M. (2008). Poor Intake of Milk, Vegetables, and Fruit with Limited Dietary Variety y Botswana's Elderly. *Journal of nutrition for the elderly*. **25**: 3–22.

Masibo, P.K. (2013). *Effects of Initial Nutritional Status on the Responses to a School Feeding Programme among School Children Aged 6 to 13 Years in the Millennium Villages Project, Siaya, Kenya*. Dissertation presented for the degree of Doctor of Philosophy (Nutritional Sciences) in the Faculty of Medicine and Health Sciences at Stellenbosch University Supervisor: Stellenbosch University.

May, P. a et al. (2014). Dietary intake, nutrition, and fetal alcohol spectrum disorders in the Western Cape Province of South Africa. *Reproductive toxicology (Elmsford, N.Y.)*,. **46C**: 31–39.

Mbochi, R.W. et al. (2012). Predictors of overweight and obesity in adult women in Nairobi Province, Kenya. *BMC public health*, **12**: 823.

Merchant, A.T. et al. (2005). Nutrient estimation from an FFQ developed for a Black Zimbabwean population. *Nutrition journal*. **4**:37.

Mitchikpe, C.E.S. et al. (2008). Seasonal variation in food pattern but not in energy and nutrient intakes of rural Beninese school-aged children. *Public health nutrition*: **12**: 414–422.

Mostert, D. et al. (2005). Dietary intake of pregnant women and their infants in a poor black South African community. *Curationis*. **28**: 12–19.

Mounir, G.M. et al. (2007). Nutritional factors affecting the menarcheal state of adolescent school girls in Alexandria. *The Journal of the Egyptian Public Health Association*, **82**: 239–260.

Msaki, M.M. and Hendriks, S.L. (2014). Measuring Household Food Security Using Food Intake Indicators in Rural Kwazulu. *Ecology of Food Nutrition*. **53**: 193–213.

Mupere, E. et al. (2012). Low nutrient intake among adult women and patients with severe tuberculosis disease in Uganda: a cross-sectional study. *BMC public health*. **12**: 1050–1050.

Nago, E.S. et al. (2010). Food, energy and macronutrient contribution of out-of-home foods in school-going adolescents in Cotonou, Benin. *The British journal of nutrition*. **103**: 281–288.

Namugumya, B.S. and Muyanja, C. (2011). Contribution of street foods to the dietary needs of street food vendors in Kampala, Jinja and Masaka districts, Uganda. *Public Health Nutrition*. **15**: 1503–1511.

Naude, C.E. et al. (2011). Growth and weight status in treatment-naïve 12-16 year old

adolescents with Alcohol Use Disorders in Cape Town, South Africa. *Nutrition journal*. **10**: 87.

Ngo, J. et al. (2009). How dietary intake methodology is adapted for use in European immigrant population groups - a review. *The British journal of nutrition*. **101 Suppl**: S86–S94.

Nutrition Coordinating Center's University of Minnesota, Nutrition Data System for Research. Available at: [www.ncc.umn.edu](http://www.ncc.umn.edu).

Nyuar, K.B., Khalil, A.K.H. and Crawford, M.A. (2012). Dietary intake of Sudanese women: a comparative assessment of nutrient intake of displaced and non-displaced women. *Nutrition and health*. **21**: 131–44.

O'Keefe, S.J.D. et al. (2007). Why do African Americans get more colon cancer than Native Africans? *The Journal of nutrition*. **137**: 175S–182S.

Ochola, S. and Masibo, P.K. (2014). Dietary intake of schoolchildren and adolescents in developing countries. *Annals of nutrition & metabolism*. **64**: 24–40.

Oguntibeju, O.O., Van der Heever, W.M.J. and Van Schalkwyk, F.E. (2007). The Interplay Between Socio-demographic Variables, Nutritional and Immune Status of HIV-positive/AIDS Patients. *Pakistan Journal of Biological Sciences*. **10**: 3592–3598.

Olayiwola, I.O. et al. (2012). Serum micronutrient status and nutrient intake of elderly Yoruba people in a slum of Ibadan, Nigeria. *Public Health Nutrition*. **17**: 455–461.

Oldewage-Theron, W. et al. (2008). Anaemia prevalence and dietary intake of elderly persons living in a peri-urban settlement in South Africa. *Journal of Family Ecology and Consumer Sciences /Tydskrif vir Gesinsekologie en Verbruikerswetenskappe*. **36**: 22–29.

Oldewage-Theron, W. et al. (2008). Health Status of an Elderly Population in Sharpeville, South Africa. *Health SA Gesondheid*, **13**: 3–17.

Oldewage-Theron, W. and Kruger, R. (2011). Dietary diversity and adequacy of women caregivers in a peri-urban informal settlement in South Africa. *Nutrition*, **27**: 420–427.

Oldewage-theron, W.H. et al. (2005). Situation analysis of an informal settlement in the Vaal Triangle. *Development Southern Africa*, **22**: 13–26.

Oldewage-Theron, W.H., Dicks, E.G. and Napier, C.E. (2006). Poverty, household food insecurity and nutrition: Coping strategies in an informal settlement in the Vaal Triangle, South Africa. *Public Health*, **120**: 795–804.

Oldewage-Theron, W.H., Kruger, R. and Egal, A.A., (2014). Socio-Economic Variables and Nutrient Adequacy of Women in the Vaal Region of South Africa. *Ecology of Food and Nutrition*, **53**: 514–527.

Oldewage-Theron, W.H., Samuel, F.O. and Djoulde, R.D. (2010). Serum concentration and dietary intake of vitamins A and E in low-income South African elderly. *Clinical Nutrition*, **29**: 119–123.

Oldewage-Theron, W.H., Samuel, F.O. and Venter, C.S. (2008). Zinc deficiency among the elderly attending a care centre in Sharpeville, South Africa. *Journal of Human Nutrition and Dietetics*, **21**: 566–574.

Ortega, R.M. et al. (2008). DIAL 1.0: Programa para evaluación de dietas y gestión de datos de alimentación. Available at: <http://www.alceingenieria.net>.

Papathakis, P.C. and Pearson, K.E. (2012). Food fortification improves the intake of all fortified nutrients, but fails to meet the estimated dietary requirements for vitamins A and B6, riboflavin and zinc, in lactating South African women. *Public Health Nutrition*. **15**: 1810–1817.

Pereko, K.K. et al. (2012). Overnutrition and associated factors among adults aged 20 years and above in fishing communities in the urban Cape Coast Metropolis, Ghana. *Public Health Nutrition*. **16**: 1–5.

Pirone, C. et al. (1993).

Nutridata Nutrition Services For The Food Industry. Available at: [www.nutridata.com](http://www.nutridata.com).

Pisa, P. et al. (2012). Social drift of cardiovascular disease risk factors in Africans from the North West Province of South Africa : the PURE study : cardiovascular topics. *Cardiovascular Journal Of Africa*. **23**: 371–378.

Pisa, P.T. et al. (2014). Inventory on the dietary assessment tools available and needed in Africa: a prerequisite for setting up a common methodological research infrastructure for nutritional surveillance, research and prevention of diet-related non-communicable diseases. *Critical Reviews in Food Science and Nutrition*.

Powell, B. et al. (2013). Wild foods from farm and forest in the East Usambara Mountains, Tanzania. *Ecology of food and nutrition*. **52**: 451–78.

Pretorius, S. et al. (2012). Feeding the emergence of advanced heart disease in Soweto : a nutritional survey of black African patients with heart failure. *Cardiovascular Journal of Africa*. **23**: 245–251.

- Rankin, D. et al. (2011). Dietary intakes assessed by 24-h recalls in peri-urban African adolescents: validity of energy intake compared with estimated energy expenditure. *European journal of clinical nutrition*. **65**: 910–919.
- Richter, L. et al. (2007). Cohort Profile : Mandela's children : The 1990 birth to twenty study in South Africa. *J Epidemiol.* **36**: 504–511.
- SAS, N. (2007). NutriLog Nutrition Software for Professionals. Available at: [www.nutrilog.com](http://www.nutrilog.com).
- Savy, M. et al. (2006). Dietary diversity scores and nutritional status of women change during the seasonal food shortage in rural Burkina Faso. *The Journal of nutrition*. **136**: 2625–32.
- Scarcella, P. et al. (2011). The impact of integrating food supplementation, nutritional education and HAART (Highly Active Antiretroviral Therapy) on the Nutritional Status of Patients living with HIV / AIDS in Mozambique : Results from the DREAM Programme. *Igiene e Sanita Pubblica - Parte Scientifica e Pratica*. **67**:41–53.
- Schap, T.E. et al. (2014). Merging dietary assessment with the adolescent lifestyle. *Journal of Human Nutrition and Dietetics*. **27**(SUPPL.1): 82–88.
- Sehmi, J. (1993). National Food Composition Tables and The Planning of Satisfactory Diets in Kenya.
- Sharma, S. et al. (1996). Development of food frequency questionnaires in three population samples of African origin from Cameroon, Jamaica and Caribbean migrants to the UK. *European Journal of Clinical Nutrition*. **50**: 479–486.
- Sheehy, T. et al. (2013). Development of a quantitative food frequency questionnaire for use

among rural South Africans in KwaZulu-Natal. *Journal of human nutrition and dietetics*. **27**: 443-449

Smith, A.F. (1991). Cognitive Processes in Long-Term Dietary Recall. **In:** *Vital Center for Health Statistics*.

Smith, A.F., Jobe, J.B. and Mingay, D.J. (1991). Retrieval from memory of dietary information. *Applied Cognitive Psychology*. **5**: 269–296.

Sodjinou, R. et al. (2009). Dietary patterns of urban adults in Benin: relationship with overall diet quality and socio-demographic characteristics. *European journal of clinical nutrition*. **63**: 222–228.

Sodjinou, R. et al. (2008). Obesity and cardio-metabolic risk factors in urban adults of Benin: relationship with socio-economic status, urbanisation, and lifestyle patterns. *BMC public health*. **8**:84.

Steyn, N. and Senekal, M. (2005). *A guide for the use of the dietary assessment and education kit (DAEK)*. Cape Town, South Africa.

Steyn, N.P. et al. (2011). Dietary, social, and environmental determinants of obesity in Kenyan women. *Scandinavian journal of public health*. **39**: 88–97.

Steyn, N.P. et al. (2012). Urbanisation and the nutrition transition: A comparison of diet and weight status of South African and Kenyan women. *Scandinavian Journal of Public Health*. **40**: 229–238.

Steyn, N.P. and Nel, J.H. (2006). Dietary Intake of Adult Women in South Africa and Nigeria

with a Focus on the Use of Spreads.

Termote, C. et al. (2012). A biodiverse rich environment does not contribute to a better diet: A case study from DR Congo. *PLoS ONE*. **7**.

Tesfaye, F., Byass, P. and Wall, S. (2008). Concurrent comparison of energy intake and expenditure among adults in Butajira District, Ethiopia. *Public health nutrition*. **11**: 675–683.

Tessier, S. et al. (2008). Regular users of supermarkets in Greater Tunis have a slightly improved diet quality. *The Journal of nutrition*. **138**: 768–774.

Thompson, F.E. and Byers, T. (1994). Dietary Assessment Resource Manual. *The Journal of Nutrition*. **124**:2245S–2317S.

Research Group Food Chemistry and Human Nutrition and Medicine, Universiteit Gent and Nutrition and Child Health Unit of the Institute of Tropical Medicine. (n.a.) Lucille food analysis software. Belgium.

US Department of Agriculture, Agricultural Research Service, N.D.L. (1996). USDA National Nutrient Database for Standard Reference, Release 28.

Venter, C.S., MacIntyre, U.E. and Vorster, H.H. (2000). The development and testing of a food portion photograph book for use in an African population. *Journal of Human Nutrition and Dietetics*. **13**: 205–218.

de Villiers, A. et al. (2006). *Relative validity of a quantified food frequency questionnaire in Black South African women*. University of Cape Town.

Vorster, H.H. et al. (2007). Cardiovascular disease risk factors and socio-economic position of



Africans in transition: the THUSA study. *Cardiovascular journal of Africa*. **18**: 282–9.

Walton, C. et al. (2012). Associations of diet quality with dairy group membership, membership duration and non-membership for Kenyan farm women and children: a comparative study. *Public health nutrition*, **17**: 1–10.

Waudu, J. et al. (2005). Food Consumption Patterns and Nutrient Intake By Women And Under Five Year Old Children In Wetlands Of Lake Victoria Basin. **In**: Food & Nutrition Security for Health and Development. Proceedings of the Inaugural National Nutrition Congress [INNC]. Nairobi, Kenya.

Wentzel-Viljoen, E., Laubscher, R. and Kruger, A. (2011). Using different approaches to assess the reproducibility of a culturally sensitive quantified food frequency questionnaire. *South African Journal of Clinical Nutrition*. **24**: 143–148.

Wiesmann, D., Arimond, M. and Loechi, C. (2009). Dietary Diversity as a Measure of the Micronutrient Adequacy of Women's Diets: Results from Rural Mozambique Site. Food And Nutrition Technic Al A Ssistance, Technical Report.

Wiig, K. and Smith, C. (2007). An Exploratory Investigation of Dietary Intake and Weight in Human Immunodeficiency Virus-Seropositive Individuals in Accra, Ghana. *Journal of the American Dietetic Association*. **107**: 1008–1013.

Willet, W. (1998). *Nutrition Epidemiology* 2nd edition. Oxford University Press, New York.

Wirfalt, E. (1998). Cognitive aspects of dietary assessment. *Scandinavian Journal of Nutrition*. **42**: 56–59.

Wojtusiak, J., Gewa, C.A. and Pawloski, L.R. (2011). Dietary Assessment in Africa: Integration with Innovative Technology. *African Journal of Food, Agriculture, Nutrition and Development*. **11**: 5629–5645.

World Health Organization. (2011). Global status report on noncommunicable diseases 2010. Available at: [http://www.who.int/nmh/publications/ncd\\_report\\_full\\_en.pdf](http://www.who.int/nmh/publications/ncd_report_full_en.pdf).

World Health Organization - Regional Office for Africa. (2006). *The African Regional Health Report: The Health of the People*. Available at: <http://www.who.int/bulletin/africanhealth/en/>.

Wrottesley, S. V et al. (2014). Europe PMC Funders Group Dietary intake and body composition in HIV-positive and - negative South African women. *Public Health Nutr.* **17**: 1603–1613.

Zeba, N.A., Delisle, H.F. and Renier G. (2014). Dietary patterns and physical inactivity, two contributing factors to the double burden of malnutrition among adults in Burkina Faso, West Africa. *Journal of Nutritional Science*. **3**: 1–14.

Zhu, F. et al. (2008). Technology-Assisted Dietary Assessment. *National Institute of Health*. **20**: 6814–681411.

Zingoni, C. et al. (2009). Studying a Population Undergoing Nutrition Transition: A Practical Case Study of Dietary Assessment in Urban South African Adolescents. *Ecology of Food and Nutrition*. **48**: 178–198.

**Table 1:** Selected studies (n=54) which assessed dietary intake of different African populations using 24-hour recall [from 2005 to 2014].

Author	Country	Study Design	Study Population	Number of recalls	Determination of Portion Sizes	Tools for Dietary Analysis
<b>2014</b>						
(Zeba et al. 2014)	Burkina Faso	Cross-sectional study	110 Adults (25-6 years-old)	2 non-consecutive	Local kitchen utensils	Malian FCT <sup>1</sup> ; C-SIDE <sup>2</sup> (Iowa State University 1996)
(Kim et al. 2014)	Tanzania	Cross-sectional study	80 Pregnant and/or Lactating Women (>18 years-old)	2 non-consecutive	Standardized food models	Tanzanian FCT Harvard University School of Public Health
(May et al. 2014)	South Africa	Case Control Study	128 Women (Mean age of 35 years-old)	Single	Photographs of local alcoholic beverage	Nutrition Data System for Research (Nutrition Coordinating Center's University of

<sup>1</sup> FCT: Food Composition Table

<sup>2</sup> C-SIDE: Software for Intake Distribution Estimation

						Minnesota n.d.)
(Changamir e et al. 2014)	Tanzania	Cohort Study	8428 Pregnant and/or Lactating Women (>18 years-old)	3	N.A. <sup>3</sup>	Tanzanian FCT Harvard University School of Public Health
(Oldewage- Theron et al. 2014)	South Africa	Cross- sectional study	722 Women (19-90 years- old)	$\geq 3$	Food models, household utensils	FoodFinder® (Grant et al. 1992)
<b>2013</b>						
(Powell et al. 2013)	Tanzania	Cross- sectional study	274 Women	2 non- consecutive	Local serving sizes aids	Programme CANDAT(Godi n 2007) ; Tanzanian FCT, FAO FTC, USDA Database (US Department of Agriculture,

<sup>3</sup> N.A.: Information not available

						Agricultural Research Service 1996) and scientific literature <sup>4</sup> .
(Koethe et al. 2013)	Zambia	Prospective Cohort study	142 Adults	4 non-consecutive	Artificial food models and serving utensils	FCT published by the Zambian National Food and Nutrition Commission; Nutrition Data System for Research (Nutrition Coordinating Center's University of Minnesota n.d.)
(Kolahdooz et al. 2013)	South Africa	Cross-sectional	137 Adults (>19 years-	2 non-consecutive	3D food models; local	Nutribase (CyberSoft

---

<sup>4</sup> Lukmanji et al. 2008; Wu Leung 1968

		study	old)		household utensils	1986)
<b>2012</b>						
(Walton et al. 2012)	Kenya	Cross-sectional study	111 Women	Single	Placing dried beans into the individual bowl to represent the serving	WorldFood Dietary Assessment System (Bunch & Murphy 1997) and the Kenyan food composition database (Sehmi 1993); USDA database <sup>5</sup> (US Department of Agriculture, Agricultural Research Service 1996)
(Termote et	Democratic	Cross-	492 Women	2 non-	1) a booklet	Lucille food

<sup>5</sup> US Department of Agriculture-Agriculture Research Service (2007) USDA National Nutrient Database for Standard Reference

al. 2012)	Republic of Congo	sectional study		consecutive	with photographs of different calibrated portion sizes; 2) an extensive price-weight- conversion list covering all foods or ingredients reported during the 24 h recalls; and 3) direct measurements of estimated leftovers	analysis software (UGent Research Group Food Chemistry and Human Nutrition & Medicine n.d.)
(Steyn et al. 2012)	Kenya	Cross- sectional study	1008 Women (15- 60 years-old)	Single	Dietary assessment kit comprising	FoodFinder® (Grant et al. 1992)

					life-size drawings and generic food models.	
(Pereko et al. 2012)	Ghana	Cross-sectional study	252 Adults (20-50)	3 non-consecutive	N.A.	ESHA Food Processor® (Davison & Mandible 1994) and the Ghanaian FCT
(López et al. 2012)	Morocco	Cross-sectional study	327 Adolescents (15-20 years-old)	3 non-consecutive	N.A.	DIAL 1.0 (Ortega et al. 2008); FCT for use in Africa, from FAO.
(Mupere et al. 2012)	Uganda	Cross-sectional study	131 Adults (>18 years-old)	Single	Local food photographs, portion-size images, volumetric vessels	East African FCT and African FCT; USDA database (US Department of Agriculture,



						Agricultural Research Service 1996); NutriSurvey Program (Erhardt & Gross 2007)
(Papathakis & Pearson 2012)	South Africa	Cohort Study	142 Pregnant and/or Lactating Women	4 non-consecutive	Volume of cups, bowls and plates	FoodFinder® (Grant et al. 1992)
(Nyuar et al. 2012)	Sudan	N.A.	113 Women (18-42 years-old)	Single	Household measures	Foodbase Nutritional Program <sup>6</sup>
(Boumaiza et al. 2012)	Tunisia	Cross-sectional study	329 Adults (Mean age of 44,9 years-old)	3 non-consecutive	Household measures	Dietetik® and Nutrilog®(SAS 2007)
<b>2011</b>						
(Steyn et al.	Kenya	Cross-	1050	Single	Photographs,	FoodFinder®

<sup>6</sup> version 4, Institute of Brain Chemistry and Human Nutrition, London Metropolitan University

2011)		sectional study	Women (15- 60 years-old)		life-size drawings food models.	(Grant et al. 1992)
(Naude et al. 2011)	South Africa	Cross- sectional study	162 Adolescents (12-16 years- old)	3 non- consecutive	Household measures;  Pictures from the Dietary Assessment and Educacioal Kit <sup>7</sup> ; MRC Food Quantities Manual.	FoodFinder® (Grant et al. 1992)
(Gibson et al. 2011)	Malawi	Cross- sectional study	80 Pregnant Women (14- 45 years-old)	3 non- consecutive	N.A.	Malawian FCT
(Hansen et al. 2011)	Kenya	Cross- sectional study	1163 Adults (18-68 years- old)	2	Real food items/Paper models,	GIES <sup>8</sup>

<sup>7</sup> Steyn and Senekal, 2002

<sup>8</sup> GIES: General Intake Estimation System Program, GIES; National Food Institute, Søborg, Denmark

					utensils from the local market	
(Addo et al. 2011)	Ghana	Cross- sectional study	70 Pregnant and/or Lactating Women (18- 42 years-old)	3 non- consecutive	Household measures, weigh portion made at home or bought	ESHA Food Processor® (Davison & Mandible 1994) and published food composition information
(Irvine et al. 2011)	Tanzania	Case Control Study	171 Adults (Mean age of 38 years-old)	Single	Real food items; Kitchen utensils; Serving dishes.	ESHA Food Processor® (Davison & Mandible 1994)
(Luke et al. 2011)	Ghana, South Africa, Seychelles, Jamaica and	Longitudinal observation study	2500 Adults (25-45 years- old)	2	Photographs; usual portions of local foods	Nutrition Data System for Research (Nutrition Coordinating

	United States					Center's University of Minnesota n.d.)
(Alemayehu et al. 2011)	Ethiopia	Cross-sectional study	68 Women (15-49 years-old)	Single	Weigh of the estimated portion consumed (using a spoon); households measures and actual food samples purchased in markets.	Ethiopian FCT and USDA database (US Department of Agriculture, Agricultural Research Service 1996)
(Rankin et al. 2011)	South Africa	Cross-sectional study	131 Adolescents	7 non-consecutive	Validated food portion photograph book (Venter et al. 2000)	FoodFinder® (Grant et al. 1992)
<b>2010</b>						

(Scarcella et al. 2011)	Mozambique	Cohort Study	106 Adults	Single	Food models and images of portion sizes	FAO FCT for Africa and Mozambique <sup>9</sup>
(Becquey & Martin-Prevel 2010)	Burkina Faso	Cross-sectional study	182 Women (19-49 years-old)	3	Household measures	C-SIDE Software (Iowa State University 1996); Malian FCT, INFOODS <sup>10</sup> database (Senegal), USDA Database (US Department of Agriculture, Agricultural Research Service 1996)
(Oldewage-Theron et	South Africa	Cross-sectional	235 Elderly (≥60 years-	2 non-consecutive:	Food Models	FoodFinder® III (Grant et al.

<sup>9</sup> Repartição de Nutrição (1991) Tabela de Composição de Alimentos Maputo:MISAU

<sup>10</sup> INFOODS: International Network of Food Data Systems

al. 2010)		study	old)	2nd 59% of the sample		1992)
(Nago et al. 2010)	Benin	Cross- sectional study	656 Adolescents (≥13 years- old)	2 non- consecutive	Household utensils.	Malian FCT; FCT for use in Africa; East African FCT.
(Dapi et al. 2010)	Cameroon	Cross- sectional study	227 Adolescents (12-16 years- old)	3 non- consecutive	Household measures, real food portions and information about the amount of money spent on some foods. Colour picture booklet	Becel Institution Nutrition Software
(Heimburge r et al. 2010)	Zambia	Cohort Study	874 Adults	4 non- consecutive	N.A.	Nutrition Data System for Research

						(Nutrition Coordinating Center's University of Minnesota n.d.)
<b>2009</b>						
(Lamri- Senhadji et al. 2009)	Algeria	Cross- sectional study	46 Adults (Mean age: 24 years-old)	5 weeks of measurement	Small, medium or large (graduated measure, soup and coffee spoons, dinner and soup plates, etc).	La composition des aliments Tableaux des valeurs nutritives*
(Sodjinou et al. 2009)	Benin	Cross- sectional study	200 Adults (25-60 years-3 old)		Local cups, bowls, spoons, plates and glasses	WorldFood Dietary Assessment System(Bunch & Murphy 1997); FCT of

						neighboring countries; C-SIDE Software (Iowa State University 1996)
(Huybregts et al. 2009)	Burkina Faso	Cross-sectional study	394 Pregnant and/or Lactating Women (15-45 years-old)	Single	Validated booklet with food photographs	Malian FCT; ESHA Food Processor® (Davison & Mandible 1994)
(Alaofè et al. 2009)	Benin	Quasi experimental	68 Adolescents (12-17 years-old)	48hR (single) 24hR (3)	Food models, portion-size models, containers and photographs of foods	Nutrifiq® software, DANA-INFRE FCT used in Benin <sup>11</sup>
(Wiesmann et al. 2009)	Mozambique	Cross-sectional study	409 Women (15-49 years-old)	Single and 2 non-consecutive	Direct weighing, volume	Specific FCT based on USDA Database (US

<sup>11</sup> DANA-INFRE: Direction de l'alimentation et de la nutrition–Institut national pour la formation et la recherche en education



					containers, photographs.	Department of Agriculture, Agricultural Research Service 1996)
(Kennedy et al. 2009)	Mali	Cross-sectional study	102 Women (15-49 years-old)	2 non consecutive	Household measures	VBS Food Calculation System (KOMEET, VBS MANAGER, ORION and FOOD GROUPS) <sup>12</sup>
(Becquey et al. 2009)	Burkina Faso	Cross-sectional study	182 Women (17-49 years-old)	3 non consecutive	Weighing of a replica, measure of the volume, use of calibrated	Malian FCT; Worldfood FCT for Senegal; USDA Database (US Department of

<sup>12</sup> Bas Nutrition Software, Arnhem, The Netherlands, [www.bware.nl](http://www.bware.nl).

					household measures, portion sizes or price	Agriculture, Agricultural Research Service 1996).
<b>2008</b>						
(Sodjinou et al. 2008)	Benin	Cross- sectional study	200 Adults (25-60 years- old)	3 non consecutive	Local cups, bowls, spoons, plates and glasses	WorldFood Dietary Assessment System (Bunch & Murphy 1997). FCT of neighboring countries; C- SIDE Software (Iowa State University 1996).
(Ijarotimi & Keshinro 2008)	Nigeria	Cross- sectional study	452 Adults (≥20 years- old)	Single	Household measures	Food analysis: AOAC <sup>13</sup> method

<sup>13</sup> AOAC: Association of Official Agricultural Chemists

(W. H. Oldewage-Theron et al. 2008)	South Africa	Cross-sectional descriptive study	170 Elderly ( $\geq 60$ years-old)	2	N.A.	FoodFinder® III (Grant et al. 1992)
(Maruapula & Chapman-Novakofski 2008)	Botswana	Cross-sectional study	99 Elderly (60-95 years-old)	Single	N.A.	Nutritionist Five <sup>14</sup>
(Gewa et al. 2008)	Kenya	Cross-sectional study	44 Women	Single	Food models, measuring cylinders, local household measures	WorldFood Dietary Assessment System (Bunch & Murphy 1997)
(Tesfaye et al. 2008)	Ethiopia	Cross-sectional study	619 Adults (18-64 years-old)	Single	Household measures, described as S, M, L., pictures of	FCT for Ethiopia

<sup>14</sup> Nutritionist Five, Version 2.3; First DataBank, San Bruno, CA 2000.

					foods and utensils	
(Oldewage-Theron et al. 2008)	South Africa	Cross-sectional study	138 Elderly (60-93 years-old)	2 non-consecutive	Food Models	FoodFinder® (Grant et al. 1992)
(WH Oldewage-Theron et al. 2008)	South Africa	Cross-sectional study	101 Elderly (60-110 years-old)	2 non-consecutive	Food Models	FoodFinder® (Grant et al. 1992)
<b>2007</b>						
(Kamau-Mbuthia & Elmadfa 2007)	Kenya	Cross-sectional study	716 Pregnant and/or Lactating Women (Reproductive age)	Single	Household measures (cups, tea and tablespoons and bowls) and also preparation methods for the different foods.	NutriSurvey Program (Erhardt & Gross 2007)

(Wiig & Smith 2007)	Ghana	Cross-sectional study	50 Adults (18-65 years-old)	Single	Food models; Portion Size images	ESHA Food Processor® (Davison & Mandible 1994) and published food composition information
(Mounir et al. 2007)	Egypt	Cross-sectional study	1606 Adolescents (Menarcheal age)	Single	N.A.	Egyptian FCT, National Nutritional Institute
(O'Keefe et al. 2007)	South Africa	Cross-sectional study	52 Adults (50-60 years-old)	3-consecutive	N.A.	South African Food Composition Database of the MRC
<b>2006</b>						
(Steyn & Nel 2006)	Kenya	Cross-sectional study	1008 Women (15-60 years-old)	Single	Household utensils, life-size drawings	Kenyan FCT and FoodFinder®

					and food models.	(Grant et al. 1992).
<b>2005</b>						
(Mostert et al. 2005)	South Africa	Cohort Study	46 Pregnant and/or Lactating Women (<40 years-old)	2	N.A.	South African Food Composition Database of the MRC
(Charlton et al. 2005)	South Africa	Cross-sectional study	285 Elderly (>60 years-old)	Single	Standard household measuring utensils, rulers and validated food photographs	FoodFinder® III (Grant et al. 1992)

**Table 2:** Selected studies (n=30) which assessed dietary intake of different African populations using food frequency questionnaires [from 2005 to 2014].

Author s	Coun try	Study Design	Study Popul ation	Dietar y Assess ment Metho d (Sourc e)	Num ber of food item s	Refere nce time frame	Validatio n/ Reprodu cibility	Determi nation of Portion Sizes	Tools for Dietary Data Analysi s
<b>2014</b>									
(Wrotte sley et al. 2014)	South Africa	Cross- section al study	247 Wome n (23- 39 years- old)	QFFQ	214	Precedi ng 7 days	N.A. <sup>15</sup>	Househo ld measures ,2D life- size drawings of foods and utensils,	FoodFi nder® III (Grant et al. 1992)

<sup>15</sup> N.A.: Information not available

								3D food models	
(Baroudi et al. 2014)	Tunisia	Case Control Study	348 Adults (20-89 years-old)	sFFQ (Decarli et al. 1996; Franceschi et al. 1993)	77	Preceding year	Tested for reproducibility and validated against 7-day WR	N.A.	Binult logiciel <sup>16</sup>
(Botha et al. 2014)	South Africa	Cross-sectional Study	1068 Adults (Mean age: 56.4 years-old)	QFFQ (MacIntyre et al. 2001b; MacIntyre et al. 2001a)	145	N.A.	Tested for reproducibility and validated against 7-day WR and biomarkers	N.A.	FoodFinder® (Grant et al. 1992)

<sup>16</sup> Binult logiciel, 2.01 version



2013									
(Sheehy et al. 2013)	South Africa	Cross-sectional study	81 Adults (19-79 years-old)	QFFQ	71	N.A.	Not tested for reproducibility or validated	Household units/3D models; weighed portions	Nutribase (CyberSoft 1986)
(Lukmanji et al. 2013)	Tanzania	Longitudinal clinical trial	1078 Pregnant and/or Lactating Women	SQFFQ	85	Preceding 3 months	Not tested for reproducibility or validated	Standard utensils and Food Models	Tanzanian FCT <sup>17</sup>
(Jordan et al. 2013)	Tanzania	Case Control Study and a validation	345 Women (26-85 years-	SQFFQ	65	N.A.	Validated against 2 non-consecutive 24hR	Household measures and solid foods in	NutriSurvey Program (Erhard

<sup>17</sup> FCT: Food Composition Table

		on study	old)					pieces or slices (in the validatio n study)	t & Gross 2007)
<b>2012</b>									
(Jackso n et al. 2012)	Botsw ana	Validat ion study	79 Adults (18-75 years- old)	QFFQ	122	Precedi ng year	Validated against 4 non- consecuti ve 24hR	Food models; househol d utensils; measurin g cups and measurin g tape.	FoodFi nder® (Grant et al. 1992)
(Pretori us et al. 2012)	South Africa	Nutriti onal Survey	50 Adults (Mean age: 47± 18	QFFQ (MacI ntyre et al. 2001b;	139	Usual intake (daily, weekly and	Tested for reproduci bility and validated	Standard ized Portions pictures and	FoodFi nder® (Grant et al. 1992)

			years-old)	MacIntyre et al. 2001a)		monthly basis)	against 7-day WR and biomarkers	utensils (cups, teaspoon s)	
(Kruger et al. 2012)	South Africa	Cross-sectional study	1325 Adults (25-64 years-old)	QFFQ (MacIntyre et al. 2001b; MacIntyre et al. 2001a)	145	N.A.	Tested for reproducibility and validated against 7-day WR and biomarkers	Food portion photograph book <sup>18</sup> ; Household measures	South African Food Composition Database MRC <sup>19</sup>
(Pisa et al.	South Africa	Cross-section	2010 Adults	QFFQ (MacI	145	N.A.	Tested for	Food models	FoodFinder®

<sup>18</sup> Venter CS, MacIntyre UE, Vorster HH. The development and testing of a food portion photograph book for use in an African population. J Hum Nutr Dietet. 2000; 13:205–18.

<sup>19</sup> Langenhoven, ML, Kruger, M, Gouws, E, Faber, M. MRC Food Composition Tables. 3rd edition. Parow Valley: Medical Research Council; 1991.; Kruger M, Sayed N, Langenhoven ML, Holing F. Composition of South African foods: vegetables and fruit. Supplement to the MRC Food Composition Tables 1991. Parow Valley: Medical Research Council; 1998; Sayed N, Frans Y, Schönfeldt HC. Composition of South African foods: milk and milk products, eggs, meat and meat products. Supplement to the MRC Food Composition Tables 1991. Parow Valley: Medical Research Council; 1999.

2012)		al study	(≥ 35 years- old)	ntyre et al. 2001b; MacInt yre et al. 2001a)			reproduci bility and validated against 7- day WR and biomarke rs	and food photogra phs	(Grant et al. 1992)
(Joffe et al. 2012)	South Africa	Cross- section al study	256 Adults (18-45 years- old)	QFFQ (Steyn & Seneka l 2005; de Villier s et al. 2006)	129	N.A.	Relative- validated	Food Photogra phs	FoodFi nder® (Grant et al. 1992)
<b>2011</b>									
(Wentz el- Viljoen et al.	South Africa	Cross- section al study	175 Adults (35-70 years-	QFFQ (MacI ntyre et al.	145	Precedi ng month	Tested for reproduci bility and	Food models, food pictures,	South African Food Compo

2011)			old)	2001b; MacInt yre et al. 2001a)			validated against 7- day WR and biomarke rs	real food, food dishes, utensils	sition Databas e _MRC, USDA databas e (US Depart ment of Agricul ture, Agricul tural Researc h Service 1996)
(Ander son et	Came roon	Cross- section	1790 Adults	QFFQ (Shar	76	Precedi ng year	N.A.	Wooden food	Several FCT <sup>20</sup>

<sup>20</sup> Tan S, Wenlock R & Buss D (1985) Immigrant Foods: Second Supplement to McCance and Widdowson's The Composition of foods. London: HMSO.; Ngosom J & Abono A (1989) Les ressources alimentaires du Cameroun: Répartition Ecologique, classification et valeur nutritive (The Food Resources of Cameroon: Ecological Distribution, Classification and Nutritional Value). Yaounde': Institut de Recherche Médicinale et d'études de

al. 2011)		al study	(24-74 years- old)	ma et al. 1996)				models and cutlery	and Microdi et Softwar e (Fletch er 1994)
(Joffe et al. 2011)	South Africa	Case Control Study	148 Wome n (18- 45 years- old)	QFFQ (Steyn & Seneka l 2005; de Villier s et al. 2006)	129	N.A.	Relative- validated	Food photogra phs	FoodFi nder® III (Grant et al. 1992)
(Aouna llah-	Tunisi a	Cross- section	1019 Adoles	SQFF Q	134	Precedi ng	Tested for	Visual tools	Tunisia n Food

plantes me'dicinales. Holland B, Welch A, Unwin I, et al. (1991) McCance and Widdowson's The Composition of Foods. London: The Royal Society of Chemistry.

Skhiri et al. 2011)		al study	cents (15-19 years- old)	(El Ati et al. 2004)		month	reproduci bility and validated		Compo sition Databas e; ESHA Food Process or® (Davis n & Mandib le 1994)
(Kruge r et al. 2011)	South Africa	Cross- section al descrip tive study	330 Adults (> 30 years- old)	QFFQ (MacI ntyre et al. 2001b; MacInt yre et	145	N.A.	Tested for reproduci bility and validated against 7- day WR	Validate d food portion photogra ph book <sup>21</sup> , common	South African Food Compo sition Databas e

<sup>21</sup> Venter CS, MacIntyre UE, Vorster HH. The development and testing of a food portion photograph book for use in an African population. J Hum Nutr Dietet. 2000; 13:205–18.

				al. 2001a)			and biomarke rs	utensils and container s	_MRC USDA databas e (US Depart ment of Agricul ture, Agricul tural Researc h Service 1996)
(Delpor t et al. 2011)	South Africa	Cross- section al study	318 Men (18-40 years- old)	SQFF Q (MacI ntyre et al. 2001a)	15 food grou ps	N.A.	Tested for reproduci bility and validated against four 24hR	N.A.	FoodFi nder® (Grant et al. 1992)



2010									
(Joffe et al. 2010)	South Africa	Case Control Study	233 Women (18-45 years-old)	QFFQ (Steyn & Seneka 2005; de Villiers et al. 2006)	129	N.A.	Relative-validated	Food Photographs	FoodFinder® III (Grant et al. 1992)
2009									
(Zingoni et al. 2009)	South Africa	Cohort Study	83 Adolescents	QFFQ (Steyn & Seneka 2005; de Villiers et al. 2006)	N.A.	Usual intake (daily, weekly and monthly basis)	Not validated or tested for reproducibility	Food Photo Manual; Food flour models; Household utensils	FoodFinder® (Grant et al. 1992)

(Goedecke et al. 2009)	South Africa	Cross-sectional study	57 Adults (18-45 years-old)	QFFQ (Steyn & Seneka 2005; de Villiers et al. 2006)	129	N.A.	Relative-validated	Food Photographs	FoodFinder® (Grant et al. 1992)
<b>2008</b>									
(Tessier et al. 2008)	Tunisia	Cross-sectional study	724 Adults	QFFQ (El Ati et al. 2004)	146	N.A.	Tested for reproducibility and validated	N.A.	ESHA Food Processor® (Davison & Mandible 1994)

(Hogenkamp et al. 2008)	South Africa	Cross-sectional study	1605 Adults (15-65 years-old)	QFFQ (95)	145	N.A.	Relative-validated	Validated photographs <sup>22</sup> ; Household measures; Food models.	FoodFinder® (Grant et al. 1992)
<b>2007</b>									
(Oguntibeju et al. 2007)	South Africa	Cross-sectional study	35 Adults (18-65 years-old)	QFFQ (MacIntyre et al. 2001b; MacIntyre et al. 2001a)	145	Preceding 6 months	Tested for reproducibility and validated against 7-day WR and biomarkers	Food models; Household measures; MRC Food Quantities Manual	FoodFinder® (Grant et al. 1992)

<sup>22</sup> Venter CS, MacIntyre UE, Vorster HH. The development and testing of a food portion photograph book for use in an African population. J Hum Nutr Dietet. 2000; 13:205–18.

(MacK eown et al. 2007)	South Africa	Cohort Study	143 Adoles cents (10 and 13 years- old)	SQFF Q (Richt er et al. 2007)	145	N.A.	Tested for Reproduc ibility and validated	NRPNI <sup>23</sup> ; Food quantitie s manual; Househo ld measures	South African Food Compo sition Databas e _MRC <sup>24</sup> ; SAS softwar e
(Vorste r et al. 2007)	South Africa	Cross- section al study	1854 Adults (≥15 years- old)	QFFQ (MacI ntyre et al. 2001b; MacInt yre et	145	N.A.	Tested for reproduci bility and validated against 7- day WR	N.A.	FoodFi nder® (Grant et al. 1992)

<sup>23</sup> National Research Programme for Nutritional Intervention

<sup>24</sup> Langenhoven, ML, Kruger, M, Gouws, E, Faber, M. MRC Food Composition Tables. 3rd edition. Parow Valley: Medical Research Council; 1991.; Kruger M, Sayed N, Langenhoven ML, Holing F. Composition of South African foods: vegetables and fruit. Supplement to the MRC Food Composition Tables 1991. Parow Valley: Medical Research Council; 1998; Sayed N, Frans Y, Schönfeldt HC. Composition of South African foods: milk and milk products, eggs, meat and meat products. Supplement to the MRC Food Composition Tables 1991. Parow Valley: Medical Research Council; 1999.

				al. 2001a)			and biomarke rs		
(Jackso n et al. 2007)	Came room	Cross- section al study	547 Adults (25-74 years- old)	QFFQ (Shar ma et al. 1996)	37	Precedi ng year	N.A.	N.A.	N.A.
<b>2006</b>									
(Belgn aoui & Belahs en 2006)	Moro cco	Cross- section al study	172 Pregna nt and/or Lactati ng Wome n (16- 44 years- old)	QFFQ	N.A.	Usual intake (daily, weekly and monthl y basis)	N.A.	N.A.	Bilnut Softwar e <sup>25</sup>

<sup>25</sup> Bilnut: SCDA Nutrisoft, Cerelles, France

(Hattin gh et al. 2006)	South Africa	Cross- section al study	488 Wome n (25- 44 years- old)	QFFQ	N.A.	N.A.	Validated	Food Quantitie s Manual	FoodFi nder® (Grant et al. 1992)
<b>2005</b>									
(Kesa & Oldewa ge- Theron 2005)	South Africa	Cross- section al study	431 Pregna nt and/or Lactati ng Wome n (16- 35 years- old)	QFFQ	N.A.	Habitu al consu mption	Validated	Food Models	Dietary Manage r Progra m®
(Merch ant et al. 2005)	Zimb abwe	Cross- section al study	100 Adults (34- 93)	SQFF Q	30	Previo us year	Validated against 24hR	Standard portion sizes	ESHA Food Process or®

									(Davison & Mandible 1994)
--	--	--	--	--	--	--	--	--	---------------------------

**Table 3:** Selected studies (n=11) which assessed dietary intake of different African populations using food frequency questionnaires and 24-hour recall [from 2005 to 2014].

Author	Country	Study Design	Study Population	Number of food items in the FFQ	Reference time frame of the FFQ	Number of recalls	Determination of Portion Sizes	Tools for Dietary Data Analysis
<b>2014</b>								
(Korkalo et al. 2014)	Mozambique	Cross-sectional study	551 Adolescents (14-19 years-old)	37	Preceding 7 days	4 non-consecutive (100%; 2 76%; 3 67%; 4 59%)	Food Photographs (Validated Food photographs in portion size estimation), Common	Mozambique FCT <sup>26</sup> for NutriSurvey Program (Erhardt & Gross 2007); Collection and

<sup>26</sup> FCT: Food Composition Table



							utensils	Analysis of foods.
<b>2012</b>								
(Mbochi et al. 2012)	Kenya	Cross - sectio nal study	365 Women (25-54 years- old)	26	Precedi ng 7 days	Single	FFQ: no used of portion sizes; 24hR:Loca l household utensils; Real foods; SA Food Photo Manual	NutriSur vey Program (Erhardt & Gross 2007)
(Amare et al. 2012)	Ethiopia	Cross - sectio nal study	356 Adults (>18 years- old)	8 food catego ries	Precedi ng 7 days	1 (100%) 3 (10%)	Household measures	Ethiopian FCT; FCT for use in Africa

(Mala et al. 2012)	Kenya	Cross-sectional study	107 Pregnant and/or Lactating Women (15-49 years-old)	N.A.	Usual intake	Single	Calibrated list of UNICEF; Household measures.	NutriSurvey Program (Erhardt & Gross 2007)
<b>2011</b>								
(Oldewage-Theron & Kruger 2011)	South Africa	N.A.	375 Women	>40	Preceding 7 days	Single	Food models	FoodFinder® (Grant et al. 1992)
(Namugumya & Muyanja 2011)	Uganda	Cross-sectional study	225 Adults (21-50 years-old)	55	Preceding week	Single	Food cost; Household utensils.	NutriSurvey Program (Erhardt & Gross 2007); USDA database

								(US Departm ent of Agricultu re, Agricultu ral Research Service 1996); FAO FCT for African foods.
<b>2010</b>								
(Baroudi et al. 2010)	Tunisia	Cross - sectio nal study	94 Adults (32-64 years- old)	168	Precedi ng month	Single	FFQ: household measures; 24hR: 3D food	Bilnut Software <sup>27</sup>

<sup>27</sup> Bilnut: SCDA Nutrisoft, Cerelles, France

							models; measurem ent aids; food specific units.	
<b>2006</b>								
(Oldewag e-Theron et al. 2006)	South Africa	Cross - sectio nal study	357 Women	N.A.	N.A.	1	Food Models	Dietary Manager Program ®
<b>2005</b>								
(Waudu et al. 2005)	Kenya, Tanzania e Uganda	Cross - sectio nal study	612 Women	N.A.	Previo us 7- day	1	Water and measuring cylinders	Food Meter UK 07
(Faber & Kruger 2005)	South Africa	Cross - sectio	187 Women (25-55	60	Precedi ng month	Single	24hR: Food models,	South African Food

		nal study	years- old)				household utensils, 3D sponge models dry oats utensils, and 3D sponge models	Composit ion Database of the MRC <sup>28</sup> .
(Oldewag e-theron et al. 2005)	South Africa	Cross sectio nal study	409 Women	N.A.	N.A.	1	Food Models	N.A.

---

<sup>28</sup> Medical Research Council

**Table 4:** Selected studies (n=4) which assessed dietary intake of different African populations using weight records [from 2005 to 2014].

Author	Country	Study Design	Study Population	Number of days	Other collected information	Tools for Dietary Data Analysis
<b>2013</b>						
(Hailelassie et al. 2013)	Ethiopia	Cross-sectional study	60 Pregnant and/or Lactating Women (15-49 years-old)	1 day	Description of the foods and their cooking methods	ESHA Food Processor® (Davison & Mandible 1994); Ethiopian FCT <sup>29</sup>
<b>2012</b>						
(Olayiwola et al. 2012)	Nigeria	Cross-sectional study	240 Elderly (>60 years-old)	3 consecutive days	Description of the what was eaten on the day before	FAO FCT; Total Dietary Assessment Software (NUTRIDATA)(Pirone & et al. 1993)

<sup>29</sup> FCT: Food Composition Table

2008						
(Gibson et al. 2008)	Ethiopia	Cross-sectional study	99 Women (Mean age: 27,8 years-old)	1 day	N.A.	Ethiopian FCT
(Abebe et al. 2008)	Ethiopia	Cross-sectional study	99 Women (Mean age: 27,8 years-old)	2 non-consecutive days	Detailed weighed recipe data for all the composite dishes	Development of a database based on the Ethiopian FCT

**Table 5:** Nutritional Tools used in dietary data analysis in the selected studies carried out between 2005-2014, organized per African Regions (Western, Eastern, Central, Northern and Southern Africa).

WESTER N AFRICA	Benin	Burkina Faso	Ghana	Mali	Nigeria
<b>Nutritional tools used in Dietary Data Analysis</b>	Nutrifiq (Alaofè et al. 2009) C-SIDE <sup>30</sup> (Sodjinou et al. 2009; Sodjinou et al. 2008) FCT <sup>31</sup> for use in Africa	USDA <sup>32</sup> Database (Becquey & Martin- Prevel 2010) ESHA Food Processor ® (Huybregts et al. 2009) C-SIDE	ESHA Food Processo r® (Addo et al. 2011; Wiig & Smith 2007; Pereko et al. 2012) Nutrition Data	VBS Food Calculation System (Kennedy et al. 2009)	Nutridata (Olayiwola et al. 2012) Food analysis methods (Ijarotimi & Keshinro 2008)

<sup>30</sup> C-SIDE: Software for Intake Distribution Estimation

<sup>31</sup> FCT: Food Composition Table

<sup>32</sup> USDA: United States Department of Agriculture



	(Nago et al. 2010)	(Zeba et al. 2014; Becquey & Martin-Prevel 2010) VBS Food Calculation System (Becquey et al. 2009)	System for Research (Luke et al. 2011)				
<b>EASTERN AFRICA</b>	<b>Ethiopia</b>	<b>Kenya</b>	<b>Tanzania</b>	<b>Uganda</b>	<b>Zambia</b>	<b>Zimbabwe</b>	<b>Mozambique</b>
<b>Nutritional tools used in Dietary Data Analysis</b>	USDA Database (Alemayehu et al. 2011) ESHA Food Processor	WorldFood Dietary Assessment System (Gewa et al. 2008; Walton et al. 2012)	NutriSurvey Program (Jordan et al. 2013) ESHA Food	USDA database, NutriSurvey Program (Mupere et al. 2012;	Nutrition Data System for Research software (Koethe et al.	USDA Database; ESHA Food Processor® (Mercha	NutriSurvey Program (Korkalo et al. 2014) FCT for Africa and

	® (Haileslassie et al. 2013) FCT for use in Africa; Ethiopian FCT (Tesfaye et al. 2008; Gibson et al. 2008; Abebe et al. 2008; Amare et al. 2012)	NutriSurvey Program (Mbochi et al. 2012; Mala et al. 2012; Kamau-Mbuthia & Elmadfa 2007) GIES <sup>33</sup> (Hansen et al. 2011) FoodFinder® (Steyn & Nel 2006; Steyn et al. 2011; Steyn et al.	Proceso r® (Irvine et al. 2011) Program me CANDA T (Powell et al. 2013) Tanzania n FCT (Kim et al. 2014; Changa mire et al. 2014; Lukmanji et al.	Namugunya & Muyanja 2011) Food Meter UK 07 (Waudonet al. 2005)	2013; Heimburger et al. 2010)	nt et al. 2005)	FCT for Mozambique (Scarcella et al. 2011)
--	---	---	---	---	-------------------------------	-----------------	--

<sup>33</sup> GIES: GIES: General Intake Estimation System Program, GIES; National Food Institute, Søborg, Denmark

		2012) Food Meter UK 07 (Waud et al. 2005)	2013) Food Meter UK 07(Waud o et al. 2005)				
<b>NORTHE RN AFRICA</b>	<b>Algeria</b>	<b>Egypt</b>	<b>Morocco</b>	<b>Sudan</b>	<b>Tunisia</b>		
<b>Nutrition al tools used in Dietary Data Analysis</b>	<i>Tableaux des valeurs nutritives</i> (Lamri- Senhadji et al. 2009)	Egyptian FCT_Nutri tion Institute (Mounir et al. 2007)	Bilnut Software (Nutrisof t) (Baroudi et al. 2010) DIAL Program me <sup>34</sup>	Foodbase Nutritional Program (Nyuar et al. 2012)	Dietetik® and Nutrilog® (Boumaiza et al. 2012) Bilnut Software (Logiciel and Nutrisoft) (Baroudi et al. 2014; Belgnaoui & Belahsen 2006) ESHA Food		

<sup>34</sup> DIAL: Programa para evaluación de dietas y gestión de datos de alimentación.

			(López et al. 2012)		Processor® (Aounallah-Skhiri et al. 2011; Tessier et al. 2008)	
<b>SOUTHERN AFRICA</b>	<b>Botswana</b>	<b>South Africa</b>		<b>CENTRAL AFRICA</b>	<b>Cameroon</b>	<b>D. R. Congo</b>
<b>Nutritional tools used in Dietary Data Analysis</b>	Nutritionist Five (Maruapula & Chapman - Novakofski 2008) FoodFinder® (Jackson et al. 2012)	FoodFinder® (Wrottesley et al. 2014; Oldewage-Theron et al. 2014; Papathakis & Pearson 2012; Pretorius et al. 2012; Joffe et al. 2011; Oldewage-Theron et al. 2010; Zingoni et al. 2009; W. H. Oldewage-Theron et al. 2008; Oguntibeju et al. 2007; Charlton et al. 2005; Rankin et al. 2011; Oldewage-		<b>Nutritional tools used in Dietary Data Analysis</b>	Microdiet (Anderson et al. 2011) Becel Software (Dapi et al. 2010)	Lucille food analysis software (Termote et al. 2012)

		Theron & Kruger 2011; Joffe et al. 2010; Faber & Kruger 2005; Wentzel-Viljoen et al. 2011; Kruger et al. 2011; Mostert et al. 2005; O’Keefe et al. 2007; Naude et al. 2011; Kruger et al. 2012; Jackson et al. 2012; Hattingh et al. 2006; Vorster et al. 2007; WH Oldewage- Theron et al. 2008; Wh Oldewage-Theron et al. 2008) Nutribase (Kolahdooz et al. 2013; Sheehy et al. 2013) Dietary Manager Program® (Kesa & Oldewage-Theron			
--	--	--	--	--	--

		2005; Oldewage-Theron et al. 2006) Nutrition Data System for Research (May et al. 2014; Luke et al. 2011) South African Food Composition Database of the MRC (Mostert et al. 2005; Kruger et al. 2011; Kruger et al. 2012; Wentzel-Viljoen et al. 2011; MacKeown et al. 2007; Faber & Kruger 2005)			
--	--	---	--	--	--

**Figure 1:** Flowchart of the paper's selection procedure.