

Cellular death of two non-*Saccharomyces* wine-related yeasts during mixed fermentations with *Saccharomyces cerevisiae*

F. Pérez-Nevado ^{b,1}, H. Albergaria ^{a,1}, T. Hogg ^c, F. Girio ^{a,*}

^a Unidade de Fisiologia Microbiana e Bioprocessos, Departamento Biotecnologia, Instituto Nacional de Engenharia Tecnologia e Inovação, Estrada do Paço do Lumiar, 22, 1649-038 Lisboa, Portugal

^b Área de Nutrición y Bromatología, Departamento de Zootecnia, Univ. de Extremadura, Badajoz, Spain

^c Escola Superior de Biotecnologia, Universidade Católica Portuguesa, Rua Dr. António Bernardino de Almeida, 4200-072 Porto, Portugal

Keywords: Non-*Saccharomyces* wine yeasts; Mixed cultures; Cellular death; Toxic compounds; Yeast–yeast interactions

Abstract

The early death of two non-*Saccharomyces* wine strains (*H. guilliermondii* and *H. uvarum*) during mixed fermentations with *S. cerevisiae* was studied under enological growth conditions. Several microvinifications were performed in synthetic grape juice, either with single non-*Saccharomyces* or with mixed *S. cerevisiae*/non-*Saccharomyces* inocula. In all mixed cultures, non-*Saccharomyces* yeasts grew together with *S. cerevisiae* during the first 1–3 days (depending on the initial inoculum concentration) and then, suddenly, non-*Saccharomyces* cells began to die off, regardless of the ethanol concentrations present. Conversely, in both non-*Saccharomyces* single cultures the number of viable cells remained high (ranging 10^7 – 10^8 CFU mL⁻¹) even when cultures reached significant ethanol concentrations (up to 60–70 g L⁻¹). Thus, at least for these yeast strains, it seems that ethanol is not the main death-inducing factor. Furthermore, mixed cultures performed with different *S. cerevisiae*/*H. guilliermondii* inoculum ratios (3:1; 1:2; 1:10; 1:100) revealed that *H. guilliermondii* death increases for higher inoculum ratios. In order to investigate if the nature of the yeast–yeast interaction was related or not with a cell–cell contact-mediated mechanism, cell-free supernatants obtained from 3 and 6 day-old mixed cultures were inoculated with *H. guilliermondii* pure cultures. Under these conditions, cells still died and much higher death rates were found for the 6 days than for the 3 day-old supernatants. This strongly indicates that one or more toxic compounds produced by *S. cerevisiae* triggers the early death of the *H. guilliermondii* cells in mixed cultures with *S. cerevisiae*. Finally, although it has not been yet possible to identify the nature of the toxic compounds involved in this phenomenon we must emphasise that the *S. cerevisiae* strain used in the present work is killer sensitive with respect to the classical killer toxins, K1, K2 and K28, whereas the *H. guilliermondii* and *H. uvarum* strains are killer neutral.

Introduction

Spontaneous wine fermentations are carried out by a succession of different yeast populations, the early stages being characterised by the growth of certain non-*Saccharomyces* yeasts belonging to the genera *Candida*, *Kloeckera* and *Hanseniaspora* and the latter stages being invariably dominated by alcohol-tolerant strains of *S. cerevisiae* (Fleet and Heard, 1993; Pretorius, 2000). The growth of some of these non-*Saccharomyces* yeasts

is nowadays considered to be beneficial to enhance the wine flavour due to higher production of important metabolites, such as glycerol, esters and higher alcohols (Romano et al., 1997; Egli et al., 1998; Granchi et al., 2002; Fleet, 2003). However, the growth of these non-*Saccharomyces* fermenting yeasts is generally limited to the first two or three days of fermentation, after which they begin to die off, giving way to strains of *S. cerevisiae* which conduct the fermentation exclusively until completion (Constantí et al., 1998; Pretorius, 2000).

The wine yeast population dynamics is generally ascribed to the higher capacity of *Saccharomyces* to withstand the changing environmental conditions of increasing ethanol and organic acids concentrations, decreasing pH and nutritional depletion

* Corresponding author. Tel.: +351 210924600; fax: +351 210924721.
E-mail address: francisco.girio@ineti.pt (F. Girio).

¹ Contributed equally to the paper.

(Pretorius, 2000). In addition, the initial microflora and the chemical composition (including fungicide and/or pesticide remains) of the grape musts, as well as extrinsic factors such as sulphur dioxide addition, fermentation temperature and the use of wine starters, exert a strong selective pressure on the survival of the yeast species along the wine fermentation process (Constanti et al., 1998; Henick-Kling et al., 1998; Ganga and Martínez, 2004). The predominant role of these classic selective pressures is currently being questioned and other, as yet undefined, microbe–microbe interactions are being put forward as potentially significant in influencing yeast successions (Ciani and Pepe, 2002; Fleet, 2003; Nissen and Arneborg, 2003).

There are several compounds produced by yeasts during alcoholic fermentations that may become inhibitory to other yeast species or strains. Aside from ethanol, certain metabolites such as short- to medium-chain fatty acids (e.g. acetic, hexanoic, octanoic and decanoic acids) can reach concentrations leading to cell death of certain yeast species, including some strains of *S. cerevisiae* (Ludovico et al., 2001; Fleet, 2003). Another inhibitory mechanism, which can occur in wine fermentations, is the so-called yeast killer activity (Shimizu, 1993; Vadasz et al., 2000; Pérez et al., 2001). This phenomenon consists in the production of specific extracellular glycoproteins by certain yeast strains (killer yeasts) that are able to kill other yeast strains (sensitive yeasts) (Woods and Bevan, 1968; Marquina et al., 2002; Schmitt and Breinig, 2002). Strains exhibiting killer-positive activity have been identified within different yeast genera associated with wine-making, including *Saccharomyces*, *Hanseniaspora*, *Pichia*, *Candida*, *Kluyveromyces*, *Zygosaccharomyces*, *Metschnikowia* and *Cryptococcus* (Chen et al., 2000; Ciani and Fatichenti, 2001).

In fermenting wine musts, oxygen and assimilable nitrogen can be rapidly depleted due to both semi-anaerobic growth conditions and poor initial nitrogen content of the grape juice. Hansen et al. (2001) reported that *T. delbrueckii* and *K. thermotolerans* are less tolerant to conditions of low available oxygen than *S. cerevisiae*. This observation is consistent with their early death during mixed fermentations with *S. cerevisiae*. On other hand, in wine fermentations where the initial microflora is mainly composed by non-*Saccharomyces* species, amino acid and vitamin consumption during the first days of fermentation can disable the subsequent growth of *S. cerevisiae* strains (Fleet, 2003).

In spite of the above-cited reports, the early death of non-*Saccharomyces* yeasts during mixed fermentations with *S. cerevisiae* strains has been scarcely described in the literature and usually is straightforward attributed to the lower ethanol tolerance of non-*Saccharomyces* by comparison to *S. cerevisiae*. Actually, only few thorough studies have been done aiming to elucidate the causes and the mechanisms underlying this phenomenon (Hansen et al., 2001; Nissen and Arneborg, 2003; Nissen et al., 2003). In the above-mentioned studies, the authors concluded that the early death of two wine-related yeasts (*K. thermotolerans* and *T. delbrueckii*) during mixed fermentations with *S. cerevisiae* was not due to the presence of ethanol or any other toxic compound but instead to a cell–cell contact-mediated mechanism. Pina et al. (2004) also reported that several *H.*

guilliermondii and *C. stellata* strains could stand much higher ethanol concentrations than previously thought. Additionally, previous work carried out with three non-*Saccharomyces* wine strains (*H. guilliermondii*, *H. uvarum* and *C. stellata*) revealed that these yeast strains are poorly fermentative due to deficient sugar uptake that is, at least in some extent, related with a nutritional limitation and not to ethanol intolerance (Albergaria et al., 2003a,b).

Aiming to clarify these findings, in the present work the early death of two non-*Saccharomyces* wine-related yeasts (*H. guilliermondii* and *H. uvarum*) under mixed cultures with *S. cerevisiae* was investigated.

Materials and methods

Yeast strains, growth media and inoculation

Microorganisms

Hanseniaspora guilliermondii NCYC 2380 (previously isolated from Douro wine region, Portugal), *H. uvarum* PYCC 4193T (Portuguese Yeast Culture Collection, UNL, Lisbon) and *Saccharomyces cerevisiae* CCMI 885 (Culture Collection of Industrial Microorganisms, INETI, Lisbon) were examined in the present study. Strains were routinely maintained at 4 °C on YEPD-agar slants (containing 20 g l⁻¹ glucose, 10 g l⁻¹ yeast extract, 20 g l⁻¹ peptone and 20 g l⁻¹ agar).

Cultivation media

Inoculum cultures were done in YEPD medium (containing 10 g l⁻¹ yeast extract, 20 g l⁻¹ peptone and 20 g l⁻¹ glucose). The single- and mixed-cultures performed with an initial sugar concentration of 200 g l⁻¹ were cultivated in a modified synthetic grape juice media (SGJ), previously described by Ciani and Ferraro (1996). Each litre of SGJ was composed of three different solutions: solution A (500 ml), solution B (250 ml), and solution C (250 ml). The final composition of SGJ was as follows (per litre): (from solution A) D-glucose, 110 g; D-fructose, 110 g; (from solution B) L-(1)-tartaric acid, 6.0 g; L-(2)-malic acid, 3.0 g; citric acid, 0.5 g; (from solution C) YNB (yeast nitrogen base without amino acids and ammonium sulfate) (Difco), 1.7 g; CAA (vitamin-free Casamino Acids) (Difco), 2.0 g; CaCl₂, 0.2 g; arginine–HCl, 0.8 g; L-(2)-proline, 1.0 g; L-(2)-tryptophan, 0.1 g. Solutions B and C were buffered at pH 3.5 with NH₄OH and H₃PO₄, respectively. The three solutions were sterilized separately at 121 °C for 20 min and then combined aseptically. At last, the medium was supplemented with 3 g l⁻¹ of both sterile yeast extract and peptone.

The mixed cultures performed with *H. guilliermondii* and *S. cerevisiae* at an initial sugar (fructose + glucose) concentration of 100 g l⁻¹ were cultivated in the same medium described above but using in solution A only half of both sugar concentration.

Inoculation cultures

In preparing the inoculum cultures of *H. guilliermondii*, *H. uvarum* and *S. cerevisiae*, cells from one YEPD-agar slant (pre-

grown for 48 h at 30 °C) were transferred to 50 ml YEPD medium in 100 ml shake flasks and incubated at 30 °C with 150 rpm of agitation for 16 h.

Single and mixed fermentations performed with 200g l⁻¹ of initial sugar

All mixed and single cultures were performed under semi-anaerobic growth conditions, in 2 l-erlenmeyers containing 1.5 l of synthetic grape juice (SGJ) at 18 °C without agitation. All single fermentations, *H. guilliermondii*, *H. uvarum* and *S. cerevisiae*, were inoculated at an initial cell concentration of 10⁶ CFU ml⁻¹. The mixed fermentation of *H. uvarum* with *S. cerevisiae* was inoculated with 7 × 10⁵ and 3 × 10⁶ CFU ml⁻¹, respectively. The mixed fermentations of *H. guilliermondii* (Hg) with *S. cerevisiae* (Sc) were inoculated with initial cell concentration of (Sc/Hg): 1.4 × 10⁶/5.4 × 10⁵ CFU ml⁻¹; 2.2 × 10⁶/4.1 × 10⁶ CFU ml⁻¹; 1.0 × 10⁴/3.8 × 10⁵ CFU ml⁻¹; 6.0 × 10¹/5.7 × 10³ CFU ml⁻¹. Samples were taken daily for viable cell counts and quantification of glucose, fructose and ethanol concentrations.

Mixed fermentations performed with 100g l⁻¹ of initial sugar

Two mixed cultures of *H. guilliermondii* (Hg) with *S. cerevisiae* (Sc) were carried out in the same growth conditions as described for the 200 g l⁻¹ single and mixed fermentations, with an initial sugar concentration of 100 g l⁻¹. One of the cultures was performed at an initial (Sc/Hg) cell ratio of 1.5 × 10⁶/2.8 × 10⁶ CFU ml⁻¹ and the other at 1.0 × 10⁵/2.9 × 10⁶ CFU ml⁻¹, respectively. Samples were taken daily for viable cell counts and quantification of glucose, fructose and ethanol concentrations.

H. guilliermondii cultivation on cell-free supernatants

Cell-free supernatants were obtained from sterile filtration (0.22 µm Millipore membranes) of one Hg single-culture, previously fermented during 8 days, and two Sc/Hg mixed cultures fermented for 3 and 6 days, respectively. Supernatants were supplemented with nitrogen source ([NH₄]₂HPO₄), vitamins, salts and yeast extract to avoid nutritional limitations. All cell-free supernatants were inoculated with 10⁶ CFU ml⁻¹ of pure Hg cultures (pre-grown for 16 h at 30 °C and 150 rpm) and cultivated in the same growth conditions used for single and mixed fermentations.

Analysis of growth

The yeasts growth during the fermentations was obtained by viable cell quantification using the classical plate count method. Samples were taken aseptically throughout the fermentations and diluted appropriately in distilled water. In the mixed fermentations, the enumeration of *H. guilliermondii* as well as *H. uvarum* cells were performed using YEPD-agar medium supplemented with 2 µg ml⁻¹ of cycloheximide (YEPD+CYH)

(Pérez et al., 2000). The number of *S. cerevisiae* viable cells in the mixed cultures was given as the difference between the total number of colonies on YEPD-agar plates and the total number of colonies on YEPD+CYH plates. YEPD-agar and YEPD+CYH plates were incubated at 30 °C for 2–4 days and enumeration was determined after no increase in colony forming units (CFU) was observed. Preliminary enumerations of single cultures of *H. guilliermondii* and *H. uvarum* were done on both YEPD-agar and YEPD+CYH medium and similar CFU numbers were obtained (data not shown). In addition, *H. guilliermondii* and *S. cerevisiae* viable cell enumeration during mixed cultures using direct plating on YEPD-agar (CFU of Hg+Sc) followed by replica-plating on YEPD+CYH plates (CFU of Hg) gave similar results to enumerations obtained from separated inoculation on YEPD-agar (CFU of Hg+Sc) and on YEPD+CYH (CFU of Hg) plates (data not shown).

Analytical assays

Cell-free samples (filtration through 0.45 µm Millipore membranes) were used to analyse different substrates and metabolites of all fermentations. Ethanol, glucose and fructose were determined by HPLC (Merck Hitachi, Darmstadt, Germany) using a Sugar-Pak™ column (Waters, Milford, USA) connected to a RI detector (L-7490, Merck Hitachi, Darmstadt, Germany). The column was eluted at 70 °C with a degassed aqueous mobile phase containing 50 mg l⁻¹ CaEDTA, at a flow rate of 0.5 ml min⁻¹. All samples were analysed in duplicate.

Killer activity strain screenings

For killer activity screening several *S. cerevisiae* reference strains for killer and sensitive traits were used: K1 killer strain (JCR2, obtained from Dr. R.B. Wickner, Section of Genetics of Simple Eukaryotes, National Institute of Diabetes, Digestive and Kidney Diseases, NIH Bethesda, MD 20892, USA); K2 killer (EX85, *S. cerevisiae* industrial wine strain isolated and selected by Regodón et al., 1997), K28 killer (GY-2-3a obtained from Manfred Schmitt, Angewandte Molekularbiologie, Universität des Saarlandes, Im Sadtwadl, Gebäude 2, D-66123 Saarbrücken, Germany); killer sensitive (EX33, *S. cerevisiae* industrial wine strain, isolated and selected by Regodón et al., 1997).

The tests were performed as described by Boone et al. (1990). Plates of MB (0.5% yeast extract, 1% peptone, 2% glucose, 2% agar, 0.003% methylene blue, 0.1 M sodium citrate) with different pH values (pH 3.5, 3.7, 4.0, 4.3, 4.7) were seeded with killer-sensitive strains (pre-grown for 48 h on YEPD-agar slants). Strains to be tested for killer activity were loaded onto the seeded agar. Colonies exhibiting clear halos on the sensitive lawns after 3–7 days of incubation at 20 °C were considered to be killer positive. To test killer-sensitive traits, MB plates were seeded with the strains to be tested and overlaid with K2, K1 and K28 killer strains. Strains exhibiting clear halos on the plates were considered to be killer sensitive towards the respective reference killer strains (K1, K2 and K28).

Results and discussion

Early death of *H. guilliermondii* and *H. uvarum* in mixed fermentations with *S. cerevisiae*

Single cultures of *H. guilliermondii*, *H. uvarum* and *S. cerevisiae* and mixed cultures of *S. cerevisiae*/*H. guilliermondii* and *S. cerevisiae*/*H. uvarum* were performed in synthetic grape juice (SGJ) under semi-anaerobic growth conditions. Viable cells, sugar consumption and ethanol production profiles of single and mixed cultures are compared in Figs. 1 and 2. The early death of *H. guilliermondii* and *H. uvarum* was observed at the beginning of both fermentations. In the mixed culture of *S. cerevisiae*/*H. guilliermondii*, both yeasts grew together during the first day of fermentation with similar maximum specific growth rates, attaining 10^7 CFU ml⁻¹ (Fig. 1-C). Then, *H. guilliermondii* cells began to die off and death was complete

(less than 1 CFU ml⁻¹) within 7–8 days of fermentation. In the mixed culture of *S. cerevisiae*/*H. uvarum*, *H. uvarum* was unable to grow and the onset of death was observed immediately after the first day of fermentation, being complete within 2–3 days (Fig. 2-C). This faster death effect was possibly due to the higher *S. cerevisiae*/*H. uvarum* inoculum ratio in comparison with the one used for *S. cerevisiae*/*guilliermondii* mixed culture. Conversely, in the single cultures of these non-*Saccharomyces* yeasts (Figs. 1-A and 2-A) the number of viable cells remained at concentrations close to 10^7 – 10^8 CFU ml⁻¹ during a much longer fermentation time (20 days in the *H. guilliermondii* culture and 15 days in the *H. uvarum* culture).

It is generally accepted that non-*Saccharomyces* yeasts begin to die off during the early stages of spontaneous wine fermentations due to their inability to tolerate the increasing ethanol concentrations present in the must medium (Egli et al., 1998; Cocolin et al., 2000; Fleet, 2003). However, recent




Fig. 1. Viable cells of *H. guilliermondii* (○) and *S. cerevisiae* (◐), total sugars (glucose + fructose) consumption (◻) and ethanol production (Δ) during single cultures of *H. guilliermondii* (A,D) and *S. cerevisiae* (B,E) and mixed culture of *H. guilliermondii* with *S. cerevisiae* (C,F).




Fig. 2. Viable cells of *H. uvarum* (◇) and *S. cerevisiae* (○), glucose and fructose consumption (□) and ethanol production (△) during single cultures of *H. uvarum* (A,D) and *S. cerevisiae* (B,E) and mixed culture of *H. uvarum* with *S. cerevisiae* (C,F).

studies have demonstrated that *H. guilliermondii* is able to stand much higher ethanol concentrations than previously thought (Pina et al., 2004). Explanations for the early death of non-*Saccharomyces* yeasts in mixed cultures with *Saccharomyces* spp., could rely upon several of factors: competition for sugar uptake, oxygen availability, nutrient limitation, presence of toxic compounds, cell–cell contact, quorum sensing (Hansen et al., 2001; Fleet, 2003; Nissen and Arneborg, 2003; Nissen et al., 2003).

The sugar and ethanol concentrations present in the *S. cerevisiae*/*H. guilliermondii* mixed fermentation at the onset of the non-*Saccharomyces* death (Fig. 1-F) were 177 and 12 g l⁻¹, respectively, and in the *S. cerevisiae*/*H. uvarum* mixed fermentation (Fig. 2-F) were 195 and 6.6 g l⁻¹, respectively. In the single cultures, however, *H. guilliermondii* and *H. uvarum* viable cell concentrations remained at high values

(ranging 10⁷–10⁸ CFU ml⁻¹) during 15–20 days of fermentation, even when ethanol had reached relatively high concentrations (60–70 g l⁻¹) (Figs. 1-D and 2-D). These results seem to indicate that the early death of non-*Saccharomyces* in mixed growth with *S. cerevisiae* was not caused by nutrient limitation or high ethanol concentrations in the culture medium. Rojas et al. (2003) found that, for mixed cultures of *H. guilliermondii* and/or *P. anomala* with *S. cerevisiae*, the onset of the non-*Saccharomyces* death always began after the initial 1–2 days of fermentation, which agrees well with our results. Moreover, Nissen and Arneborg (2003) reported similar results for two non-*Saccharomyces* wine yeasts (*Kluyveromyces thermotolerans* and *Torulospora debreueckii*) in mixed cultures with *S. cerevisiae*. These authors suggested that viable *S. cerevisiae* cells at high concentrations provoke the early death of non-*Saccharomyces*

yeasts in mixed cultures by a cell-to-cell contact-mediated mechanism.

In the current study, the early death of *H. guilliermondii* and *H. uvarum* yeasts in mixed growth with *S. cerevisiae* was also confirmed under different cultivation conditions, namely

aerobic growth and different culture media, and tested against several *S. cerevisiae* strains (data not shown). These findings led us to conclude that this early death is not dependent of the growth conditions or of a specific *S. cerevisiae* strains but it is rather related to the physical presence of *S. cerevisiae* cells and/




Fig. 3. Viable cells of *H. guilliermondii* (Hg) (○) and *S. cerevisiae* (Sc) (◻), glucose and fructose consumption (◻) and ethanol production (Δ), during the mixed cultures performed with different (Sc/Hg) inoculum ratios: A, A'—(3:1); B, B'—(1:2); C, C'—(1:10); D, D'—(1:100).

Table 1
Relation between the initial Sc/Hg inoculum ratio and the onset of the Hg cell death in each Sc/Hg mixed culture plotted in Fig. 3

Plots in Fig. 3	Initial cell density (CFU ml ⁻¹)		Initial day of Hg cell death	<i>S. cerevisiae</i> cell density at initial day of death (CFU ml ⁻¹)
	<i>S. cerevisiae</i>	<i>H. guilliermondii</i>		
A	1.4×10^6	5.4×10^5	1	6.6×10^6
B	2.2×10^6	4.1×10^6	2	5.1×10^6
C	1.0×10^4	3.8×10^5	3	6.9×10^6
D	6.0×10^1	5.7×10^3	15	3.6×10^6

or toxic agents produced by them against other non-*Saccharomyces* yeasts.

.Effect of *S. cerevisiae* viable cells concentration on the cell death process

In order to evaluate the influence of *S. cerevisiae* cell density on the non-*Saccharomyces* death process, several mixed cultures were performed using different *S. cerevisiae*/*H. guilliermondii* (Sc/Hg) inoculum ratios (3 : 1); (1 : 2); (1 : 10); (1 : 100).

Viable cell concentrations of *H. guilliermondii* and *S. cerevisiae* of these mixed cultures and the corresponding sugars and ethanol profiles, throughout the cultivation period, are represented on Fig. 3. Plots A–D show the relationship between death kinetics and initial Sc/Hg cell concentration ratio. In all mixed cultures, *Saccharomyces* became dominant at the end of the fermentation but *H. guilliermondii* death began, and was

completed, at varying stages of the fermentation depending on the initial Sc/Hg cell concentration ratio (Fig. 3). Results show that death rates of *H. guilliermondii* in the mixed cultures increased for higher Sc/Hg inoculum ratios and death process began sooner: at the first day of fermentation for the initial inoculum ratio of (3 : 1) and after 15 days for the initial inoculum ratio of (1 : 100).

Comparing *S. cerevisiae* with *H. guilliermondii* cell profiles (Fig. 3, plots A–D) it can be observed that the onset of the non-*Saccharomyces* death depends, not only from the initial relative proportions of *S. cerevisiae*/*H. guilliermondii* but also from the *S. cerevisiae* cell density value present in the mixed cultures. In fact, *H. guilliermondii* death in the mixed cultures occurred only when *S. cerevisiae* cell density reached values close to 10^7 CFU ml⁻¹. The initial number of viable cells, the initial day of *H. guilliermondii* death process and the corresponding *S. cerevisiae* cell density in the mixed cultures are summarised in Table 1.

Once again, it is clear that the sugars and ethanol concentrations were not the primary cause of *H. guilliermondii* death (Fig. 3, plots A'–D'). In fact, at the onset of Hg death the sugars and ethanol concentrations were 187 g l⁻¹ and 0 g l⁻¹, respectively, in the (3 : 1) Sc/Hg mixed culture (Fig. 3-A') and 52 g l⁻¹ and 72 g l⁻¹, respectively, in the (1 : 100) Sc/Hg mixed culture (Fig. 3-D'). These results prove that the early death of *H. guilliermondii* is highly dependent of the *S. cerevisiae* concentration and two explanations suggest themselves: production of toxic compounds, other than ethanol, by *S. cerevisiae*, or cell-




Fig. 4. Viable cells of *H. guilliermondii* (○) and *S. cerevisiae* (□), glucose and fructose consumption (□) and ethanol production (△) during the mixed cultures performed with 100 g l⁻¹ of initial total sugars at different (Sc/Hg) inoculums ratios: A, A'—(1 : 2); B, B'—(1 : 10).

to-cell contact, as previously reported by Nissen et al. (2003) and Nissen and Arneborg (2003).

The amount of sugar metabolised by *S. cerevisiae* had a pronounced effect on the death rate of *H. guilliermondii* in mixed cultures. In fact, for the mixed culture performed with 100 g l⁻¹ of initial sugars (Sc/Hg inoculum ratio ≈1:10), the viable cells concentrations of *H. guilliermondii* remained at values close to 10⁸ CFU ml⁻¹ until the 7th day of fermentation and death was not completed, at least, during the first 34 days of fermentation (Fig. 4-B). Whereas in the mixed culture performed with 200 g l⁻¹ of initial sugars (similar inoculum ratio) *H. guilliermondii* death began on the 4th day of fermentation and was completed within 12–13 days (Fig. 3-C). A similar delay on the onset of cellular death was observed when a mixed culture was grown at (1:2) inoculum ratio (Sc/Hg) using 100 g l⁻¹ instead of 200 g l⁻¹ of initial total sugars (Fig. 4-A compares with Fig. 3-B), although *S. cerevisiae* caused the total death of *H. guilliermondii* cells in both situations. Cell densities achieved by *S. cerevisiae* in the mixed fermentations performed with 100 g l⁻¹ of initial sugars were similar to those found in the mixed fermentations performed with 200 g l⁻¹ (above 10⁷ CFU ml⁻¹), but *H. guilliermondii* death was retarded in the former case when compared to the latter. Thus, apparently, death of *H. guilliermondii* depends not only on the *S. cerevisiae* cell density but also on the amount of sugar metabolised. These findings suggest that one or more toxic compounds produced by *S. cerevisiae* is/are the cause of the early death of *H. guilliermondii* during mixed growths with *S. cerevisiae*.

It should be mentioned that the overall behaviour of the different yeast species in the mixed fermentations performed agree well with the yeast population dynamics found in spontaneous and inoculated wine fermentations. Several studies (Constanti et al., 1998; Cocolin et al., 2000; Raspor et al., 2002) report the dominance of non-*Saccharomyces* yeasts during the firsts days (3–7 days) of spontaneous vinifications, where the initial yeast population ranges 10⁴–10⁵ CFU ml⁻¹ for non-*Saccharomyces* yeasts and 10²–10³ CFU ml⁻¹ for *Saccharomyces* yeasts. This performance is similar to the yeast behaviour exhibited in the mixed fermentations represented in Fig. 3-C and D. Conversely, in vinifications where *S. cerevisiae* starters




Fig. 5. Viable cells profiles of *H. guilliermondii* inoculated in free-cell supernatants obtained from previous 8 day-old Hg single-culture (○), 3 day-old Sc/Hg mixed culture (○●) and 6 day-old Sc/Hg mixed culture (●).

Table 2

Killer activity tests performed for *S. cerevisiae* CCM1 885, *H. guilliermondii* NCYC 2380 and *H. uvarum* PYCC 4193T with the reference strains, K1 killer JCR2, K2 killer EX85, K28 killer GY-2-3a and killer-sensitive EX33

Strains	Killer activity at different pH				
	3.5	3.7	4.0	4.3	4.7
<i>H. guilliermondii</i>	N1	N1	N1	N1	N1
	N2	N2	N2	N2	N2
	N28	N28	N28	N28	N28
	N33	N33	N33	N33	N33
<i>H. uvarum</i>	N1	N1	N1	N1	N1
	N2	N2	N2	N2	N2
	N28	N28	N28	N28	N28
	N33	N33	N33	N33	N33
<i>S. cerevisiae</i>	S1	S1	S1	S1	S1
	S2	S2	S2	S2	S2
	S28	S28	S28	S28	S28
	N33	N33	N33	N33	N33

N1: Killer neutral with respect to K1 killer toxin; N2: Killer neutral with respect to K2 killer toxin; N28: Killer neutral with respect to K28 killer toxin; N33: Killer neutral with respect to killer-sensitive strain EX33; S1: Killer sensitive with respect to K1 killer toxin; S2: Killer sensitive with respect to K2 killer toxin; S28: Killer sensitive with respect to K28 killer toxin.

are added to the musts at an initial cell concentration of 10⁶ CFU ml⁻¹, the *S. cerevisiae* dominance is established within the firsts 1–2 days, which compares well with the results obtained in the mixed cultures A and B (Fig. 3-A,B).

Presence of toxic compounds

In order to investigate the cause of the non-*Saccharomyces* early death, cell-free supernatants obtained from 3 and 6 days-old mixed cultures (Hg+Sc) and from an 8 days-old single culture of Hg were inoculated with *H. guilliermondii* at an initial cell density of 10⁶ CFU ml⁻¹. The viable cell profiles of these cultures (Fig. 5) showed that total death of *H. guilliermondii* was achieved within 5 and 13 days when cultivated in the 6 and 3 day-old mixed culture supernatants, respectively. Conversely, when cultivated in the Hg single-culture supernatant, cell viability remained relatively stable during 13 days, with a slight decrease being observed from an initial cell density of 2 × 10⁶ CFU ml⁻¹ to a final cell density of 8 × 10⁴ CFU ml⁻¹. The initial content of sugars and ethanol in the mixed cultures supernatants were 138 and 26 g l⁻¹, respectively, in the 3 day-old supernatant and 74 and 58 g l⁻¹, respectively, in the 6 day-old supernatant. It should be pointed out that the latter values compare with those initially present on




Fig. 6. Pictures of the killer activity plate tests performed for *S. cerevisiae* CCM1 885 (Sc), *H. guilliermondii* NCYC 2380 (Hg) and *H. uvarum* PYCC 4193T (Hu) with the reference killer strain (K1).

the Hg single-culture supernatant (initial concentration of sugars and ethanol of 67 and 52 g l⁻¹, respectively). These results sustain the conclusion that the primary cause of the early death of *H. guilliermondii* in mixed growth with *S. cerevisiae* is not ethanol or nutrient depletion (all cell-free supernatants were supplemented with nitrogen and yeast extract). Conversely to the results reported by Nissen et al. (2003), where the early death of *K. thermotolerans* and *Torulospira delbrueckii* in mixed cultures with *S. cerevisiae* suggested a cell–cell contact-mediated mechanism, our results strongly indicate that some toxic compounds produced exclusively by *S. cerevisiae* are responsible for the early death of *H. guilliermondii* yeast.

It is well known that during wine fermentations yeasts can produce, besides ethanol, other toxic compounds, namely, killer toxins, short- and medium-chain fatty acids and sulphite, able to induce death of other yeasts — *Saccharomyces* spp. or non-*Saccharomyces* spp. (Schmitt and Breinig, 2002; Fleet, 2003). Several killer strains of *S. cerevisiae*, isolated from wine musts, have been found to kill other sensitive yeasts within the firsts days of fermentation, although the majority of these toxins are only effective against other *S. cerevisiae* strains (Shimizu, 1993; Pérez et al., 2001). To assure that classical killer toxins produced by *S. cerevisiae* were not involved in this early death phenomenon, killer activity of the three yeast strains used in the present work was tested against the classical killer toxins K1, K2, and K28 (at different pH). Tests revealed that *S. cerevisiae* CCMI 885 is killer-sensitive against the reference killer toxins (K1, K2 and K28) and *H. guilliermondii* NCYC 2380 as well as *H. uvarum* PYCC 4193T are killer neutral (Table 2 and Fig. 6). Additionally, none of the strains is killer positive towards the killer-sensitive strain EX33 (Table 2). Although significant, these results do not exclude however the production by *S. cerevisiae* of any other unknown killer-like toxins (e.g. peptides, proteins or glycoproteins) or even some fermentative metabolites potentially toxic for some non-*Saccharomyces* strains. To determine the nature of the toxic compounds involved in the early death of these non-*Saccharomyces* strains, as well as the death mechanism underlying this phenomenon, further investigation will be carried out.

Acknowledgements

The authors would like to thank Dr. J. C. Rivas for supplying yeast killer strain K28. The present work was supported by project POCTI/AGR/39974/2001, funded by Fundação para a Ciência e Tecnologia. Author F. Pérez-Nevado was beneficiary of a post-doc grant supported by Consejería de Educación, Ciencia y Tecnología (Junta de Extremadura, Spain) and European Social Fund.

References

Albergaria, H., Torão, A.R., Hogg, T., Gírio, F., 2003a. Physiological behaviour of *Hanseniaspora guilliermondii* under aerobic glucose-limited continuous cultures. *FEMS Yeast Research* 3, 211–216.

Albergaria, H., Torão, A.R., Hogg, T., Gírio, F., 2003b. The growth kinetics and fermentation behaviour of some non-*Saccharomyces* wine yeasts: *Hanse-*

niaspora uvarum, *H. guilliermondii* and *Candida stellata*. 23rd ISSY, Budapest, Hungary.

Boone, C., Sdicu, A.M., Wagner, J., Degre, R., Sánchez, C., Bussey, H., 1990. Integration of the yeast K1 killer toxin gene into the genome of marked wine yeasts and its effect on vinification. *American Journal of Enology and Viticulture* 41, 37–42.

Chen, W.B., Han, Y.F., Jong, S.C., Chang, S.C., 2000. Isolation, purification, and characterization of a killer protein from *Schwanniomyces occidentalis*. *Applied and Environmental Microbiology* 66, 5348–5352.

Ciani, M., Faticenti, F., 2001. Killer toxin of *Kluyveromyces phaffii* DBVPG 6076 as a biopreservative agent to control apiculate wine yeasts. *Applied and Environmental Microbiology* 67, 3058–3063.

Ciani, M., Ferraro, L., 1996. Enhanced glycerol content in wines made with immobilized *Candida stellata* cells. *Applied and Environmental Microbiology* 62, 128–132.

Ciani, M., Pepe, V., 2002. The influence of pre-fermentative practices on the dominance of inoculated yeast starter under industrial conditions. *Journal of the Science of Food and Agriculture* 82, 573–578.

Cocolin, L., Bisson, L.F., Mills, D.A., 2000. Direct profiling of the yeast dynamics in wine fermentations. *FEMS Microbiology Letters* 189, 81–87.

Constanti, M., Reguant, C., Poblet, M., Zamora, F., Mas, A., Guillamon, J.M., 1998. Molecular analysis of yeast population dynamics: effect of sulphur dioxide and inoculum on must fermentation. *International Journal of Food Microbiology* 41, 169–175.

Egli, C.M., Edinger, W.D., Mitrakul, C.M., Henick-Kling, T., 1998. Dynamics of indigenous and inoculated yeast populations and their effect on the sensory character of Riesling and Chardonnay wines. *Journal of Applied Microbiology* 85, 779–789.

Fleet, G.H., 2003. Yeast interactions and wine flavour. *International Journal of Food Microbiology* 86, 11–22.

Fleet, G.H., Heard, G.M., 1993. Yeasts: growth during fermentation. In: Fleet, G.H. (Ed.), *Wine Microbiology and Biotechnology*. Harwood Academic Publishers, Chur, Switzerland, pp. 27–54.

Ganga, M.A., Martínez, C., 2004. Effect of wine yeast monoculture practice on the biodiversity of non-*Saccharomyces* yeasts. *Journal of Applied Microbiology* 96, 76–83.

Granchi, L., Ganucci, D., Messini, A., Vincenzini, M., 2002. Oenological properties of *Hanseniaspora osmophila* and *Kloeckera corticis* from wines produced by spontaneous fermentations of normal and dried grapes. *FEMS Yeast Research* 2, 403–407.

Hansen, E.H., Nissen, P., Sommer, P., Nielsen, J.C., Arneborg, N., 2001. The effect of oxygen on the survival of non-*Saccharomyces* yeasts during mixed culture fermentations of grape juice with *Saccharomyces cerevisiae*. *Journal of Applied Microbiology* 91, 541–547.

Henick-Kling, T., Edinger, W., Daniel, P., Monk, P., 1998. Selective effects of sulfur dioxide and yeast starter culture addition on indigenous yeast populations and sensory characteristics of wine. *Journal of Applied Microbiology* 84, 865–876.

Ludovico, P., Sousa, M.J., Silva, M.T., Leao, C., Corte-Real, M., 2001. *Saccharomyces cerevisiae* commits to a programmed cell death process in response to acetic acid. *Microbiology* 147, 2409–2415.

Marquina, D., Santos, A., Peinado, J.M., 2002. Biology of killer yeasts. *International Microbiology* 5, 65–71.

Nissen, P., Arneborg, N., 2003. Characterization of early deaths of non-*Saccharomyces* yeasts in mixed cultures with *Saccharomyces cerevisiae*. *Archives of Microbiology* 180, 257–263.

Nissen, P., Nielsen, D., Arneborg, N., 2003. Viable *Saccharomyces cerevisiae* cells at high concentrations cause early growth arrest of non-*Saccharomyces* yeasts in mixed cultures by a cell–cell contact-mediated mechanism. *Yeast* 20, 331–341.

Pérez, F., Regodón, J.A., Valdés, M.E., De Miguel, C., Ramírez, M., 2000. Cycloheximide resistance as marker for monitoring yeasts in wine fermentations. *Food Microbiology* 17, 119–128.

Pérez, F., Ramírez, M., Regodón, J.A., 2001. Influence of killer strains of *Saccharomyces cerevisiae* on wine fermentation. *Antonie van Leeuwenhoek* 79, 393–399.

Pina, C., Santos, C., Couto, J.A., Hogg, T., 2004. Ethanol tolerance of five non-*Saccharomyces* wine yeasts in comparison with a strain of *Saccharomyces*

- cerevisiae* — influence of different culture conditions. Food Microbiology 21, 439–447.
- Pretorius, I.S., 2000. Tailoring wine yeast for the new millennium: novel approaches to the ancient art of winemaking. Yeast 16, 675–729.
- Raspor, P., Cus, F., Jemec, K.P., Zagorc, T., Cadez, N., Nemanic, J., 2002. Yeast population dynamics in spontaneous and inoculated alcoholic fermentations of Zametovka must. Food Technology and Biotechnology 40, 95–102.
- Regodón, J.A., Pérez, F., Valdés, M.E., De Miguel, C., Ramírez, M., 1997. A simple and effective procedure for selection of wine yeast strains. Food Microbiology 14, 247–254.
- Rojas, V., Gil, J.V., Pinaga, F., Manzanares, P., 2003. Acetate ester formation in wine by mixed cultures in laboratory fermentations. International Journal of Food Microbiology 86, 181–188.
- Romano, P., Suzzi, G., Comi, G., Zironi, R., Maifreni, M., 1997. Glycerol and other fermentation products of apiculate wine yeasts. Journal of Applied Microbiology 82, 615–618.
- Schmitt, M.J., Breinig, F., 2002. The viral killer system in yeast: from molecular biology to application. FEMS Microbiology Review 26, 257–276.
- Shimizu, K., 1993. Killer yeast. In: Fleet, G.H. (Ed.), Wine Microbiology and Biotechnology. Harwood Academic Publishers, Chur, Switzerland, pp. 243–263.
- Vadasz, A.S., Jagganath, D.B., Pretorius, I.S., Gupthar, A.S., 2000. Electron microscopy of the K-2 killer effect of *Saccharomyces cerevisiae* T206 on a mesophilic wine yeast. Antonie van Leeuwenhoek International Journal of General and Molecular Microbiology 78, 117–122.
- Woods, D.R., Bevan, E.A., 1968. Studies on the nature of the killer factor produced by *Saccharomyces cerevisiae*. Journal of General Microbiology 51, 115–126.