




Growth and Micropropagation Assays of *Salicornia* and *Sarcocornia* Plants



de J.-Raposo, MF; Oliveira, SA and Morais, RM*

Escola Superior de Biotecnologia – Universidade Católica Portuguesa

R. Dr. António Bernardino de Almeida, 4200 – 072 Porto

*e-mail: rmorais@esb.ucp.pt

INTRODUCTION

Salicornia and *Sarcocornia* are two succulent, halophytic plants used in salads or as pickles, hence the name “pickleweed”. Their seeds are very rich in edible oils, highly polyunsaturated, linoleic acid contributing with 70% to the fatty acids content. *Salicornias* also have high amounts of iodine, phosphorus, zinc, and vitamins A, C and D.

Due to the increasing interest in the food industry and also because it is urgent to replace these plants in their habitats, it is also natural that one thinks in their quick and efficient propagation.

At a laboratorial level, micropropagation is the process used to obtain plants with the same phenotypic and genetic characteristics, usually involving manipulation of the culture media, adding (or not) some growth regulators, to allow development of stems or roots of the cultivated explants.





Figure 1. Details of the germination of *Salicornia*/ *Sarcocornia* seeds.

METHODOLOGIES

Seeds disinfection

- ✓ ethyl alcohol 70%
- ✓ sterilised deionised water
- ✓ sodium hypochlorite 2.5% + 0.5% tween 20
- ✓ sterilised deionised water, 3x
- ✓ benomil solution 1%




Figure 2. Another aspect of the germination: when adding the growth regulators.

Germination

- ☞ in Petri dishes, with H&A medium (fig 1A) or in tubes (fig 1B),
- ☞ embedded (or not) in growth regulators GA3, 100mg l⁻¹, or salicylic acid, 7mg l⁻¹ (fig 2),
- ☞ put to germinate in media without hormones or with 0.1mg l⁻¹ IAA + 0.5mg l⁻¹ BAP (Mei *et al.*, 1997), or 2mg l⁻¹ NAA + 1mg l⁻¹ BAP + 1 GA3 (Libik *et al.*, 2005)



Figure 3. Picture represents plantlets growing in tubes and in glass flasks, in H&A medium

Growth of plantlets

- ◊ in tubes (fig 3A) or in small glass flasks (fig 3B)
- ◊ in (2)H&A medium (for a better growth)
- ◊ growth chamber, under constant temperature (25°C) and illumination (29.18 μEs⁻¹m⁻²)

Induction of organogenesis

Stem	1 st assay	IBA, NAA, 2,4-D 0.025mg l ⁻¹
	kin	0.5, 1.0, 1.5, 2.0 mg l ⁻¹ (fig 4A)
2 nd assay	IAA, IBA, 2,4-D	0.5, 1.0 mg l ⁻¹
	kin	0.5, 1.0, 1.5, 2.0 mg l ⁻¹ (fig 4B)
Root	1 st assay	IBA, 2,4-D 0.625 mg l ⁻¹
	kin	0.037 mg l ⁻¹ (fig 5A)
2 nd assay	IBA	1.825 mg l ⁻¹ (fig 5B)



Figure 4. Organogenesis assays with *Salicornia* explants to induce stem growth.

Figure 5. Organogenesis assays with *Salicornia* to induce root growth.

When nitrogen and phosphorus were increased plant answer was positive and plantlets begin to show a better growth (fig 6A and B).





Figure 6. Salicornias show better growth when in doubled concentration of nitrogen and phosphorus.

Explants are also responding well to the addition of casein hydrolysate (HC) (100 and 200 mg l⁻¹), to which BAP (benzyl aminopurine) was added, best growth obtained with 200 mg l⁻¹ HC plus 1 mg l⁻¹ BAP.

A good answer was also obtained when using explants with 2-3 segments, either in a supplemented medium or not (fig 7).

Figure 7. Explants of *Salicornia* with 1-3 segments, one week (A) and two weeks (B) after being cultivated in 2H&A medium with HC and vitamins.



- REFERENCES
- Ahmad, N; Anis, M (2005) *In vitro* Mass Propagation of *Cucumis sativus* L. from Nodal Segments. *Turk J. Bot.*, **29**: 237-240
 - Libik, M; Konieczny, R; Pater, B (2005) Differences in the Activities of Some Antioxidant Enzymes and H₂O₂ Content During Rhizogenesis and Somatic Embryogenesis in Callus Cultures of the Ice Plant. *Plant Cell Rep.*, **23**: 834-841
 - Mei, B; No, EG; McWilliams, EL; Gould, JH; Newton, RJ (1997) *In vitro* Regeneration of Fourwing Salt Bush *Atriplex canescens* (Pursh) Nutt. *Journal of Range Management*, **50**(4): 413-418
 - Prehn, D; Serrano, C; Berrios, CG; Arce-Johnson, P (2003) Micropropagation of *Quillaja saponaria* Mol. Starting from seed. *Bosque*, **24**(2): 3-12