

Workshop

Predictive Microbiology

Aplicações e Novas Fronteiras

Modelos para inactivação de microrganismos em alimentos

Teresa Brandão

Cristina Silva



UNIVERSIDADE CATÓLICA PORTUGUESA
Escola Superior de Biotecnologia

4 de Março 2005


Modelagem matemática

alguns conceitos ...

$$y = f(x, q) + e$$

Estimativa dos parâmetros

minimização dos desvios entre os valores **experimentais** e os obtidos pelo **modelo**



Precisas ?

q^*

Exactas ?



Modelagem matemática

objectivo

descrição **precisa** e **exacta** das observações

adequabilidade do modelo

qualidade dos parâmetros



Modelagem matemática

modelos mecanísticos

- ◆ descrição fundamental dos processos físico-químicos envolvidos
- ◆ mais complexos

modelos empíricos

- ◆ *caixa preta*
- ◆ mais simples (ou não!)
- ◆ aplicação prática



ponderar **vantagens** e **desvantagens**
decisão depende do **objectivo final**

Modelagem matemática

complexidade
matemática



descrição adequada

qualidade

modelo



parâmetros

vantagens

- conhecimento do processo
- efeito do processo no produto
- controlo das variáveis envolvidas


Modelagem matemática

Fenómenos de Transferência


- calor
- massa
- *momentum*

Cinéticas de reacção

Propriedades



Modelagem




Modelagem matemática

Fenómenos de Transferência

- calor
- massa
- *momentum*

Cinéticas de reacção

Propriedades





Modelagem matemática

Fenómenos de Transferência


- calor
- massa
- *momentum*

Cinéticas de reacção

Propriedades



Modelagem





Modelagem matemática

Fenómenos de Transferência


- calor
- massa
- *momentum*

Cinéticas de reacção

Propriedades



Modelagem



Modelagem matemática

Fenómenos de Transferência

- calor
- massa
- *momentum*

Cinéticas de reacção

Propriedades

Processos

Físicos

Químicos

Processos Alimentares

Modelagem

dados

design experimental

Função matemática

Esquemas de regressão

parâmetros

variáveis

design


Critério

validação

controlo

optimização

Objectivos





Modelagem matemática

Fenómenos de Transferência





- calor
- massa
- *momentum*

Cinéticas de reacção

Propriedades



Modelagem





segurança


A red starburst graphic with multiple points, centered on the page. The text 'microbiologia predictiva' is written in white, bold, sans-serif font inside the starburst.

**microbiologia
predictiva**

Modelagem matemática

**microbiologia
predictiva**

microbiologia



estatística

matemática

aplicação

- previsão / simulação → rapidez
- desenvolvimento de processos eficazes de inativação




contribuição para **segurança**

inactivação




comportamento sigmoidal




reflecte a presença de *agregados* de microrganismos
ou sub populações mais **resistentes** à **temperatura**
(ou outro **factor adverso**)

Exemplos



Miller (2004)

Exemplos




Miller (2004)

Modelos matemáticos

❖ primários




parâmetros



Modelos matemáticos


❖ primários



❖ secundários
parâmetros

Modelos matemáticos

❖ primários




❖ secundários
parâmetros

❖ terciários - integração dos modelos anteriores - **software**

Modelos de inativação

❖ primários




empíricos

fundamentais

Modelos de inativação

❖ primários



$$N = N_0 \exp(-kt)$$




$$\log N = \log N_0 - \frac{t}{D}$$

1ª ordem

D – tempo de redução decimal

Modelos de inactivação

❖ primários



$$N = N_0 \exp(-kt)$$



$$\log N = \log N_0 - \frac{t}{D}$$

1ª ordem


D – tempo de redução decimal

$$\frac{N}{N_0} = F_1 \exp(-k_1 t) + (1 - F_1) \exp(-k_2 t)$$

Cerf
(1977)

bifásicos

F_1 – fracção de microrganismos inactivados
 k_1 e k_2 – constantes cinéticas




$$\log \frac{N}{N_0} = \log \left(\frac{2F_1}{1 + \exp(k_1 t)} + \frac{2(1 - F_1)}{1 + \exp(k_2 t)} \right)$$

Kamau et al.
(1990)

Modelos de inativação

❖ primários




Whiting & Buchanan
(1992)

$$\log \frac{N}{N_0} = \log \left(\frac{F_1(1 + \exp(-k_1L))}{1 + \exp(k_1(t - L))} + \frac{(1 - F_1)(1 + \exp(-k_2L))}{1 + \exp(k_2(t - L))} \right)$$

L – atraso

Modelos de inativação

❖ primários



Whiting & Buchanan
(1992)

$$\log \frac{N}{N_0} = \log \left(\frac{F_1(1 + \exp(-k_1L))}{1 + \exp(k_1(t - L))} + \frac{(1 - F_1)(1 + \exp(-k_2L))}{1 + \exp(k_2(t - L))} \right)$$

L – atraso


Cole et al.
(1993)

$$\log N = \alpha + \frac{w - \alpha}{1 + \exp\left(\frac{4\sigma(\lambda - \log t)}{w - \sigma}\right)}$$

Distribuição da sensibilidade ao calor da população dos microrganismos

Modelos de inativação

❖ primários



Baranyi et al.
(1993)

$$\frac{dN}{dt} = -k \alpha(t) \beta(t) N$$


$$N(t = 0) = N_0$$

função 'cauda'

função 'atraso'

Modelos de inativação

❖ primários



Baranyi et al.
(1993)

$$\frac{dN}{dt} = -k \alpha(t) \beta(t) N$$

$$N(t = 0) = N_0$$

função 'cauda'

função 'atraso'

Geeraerd et al.
(2000)

$$\frac{dN}{dt} = -k_{\max} k_Q(Q) N$$


$$\frac{dQ}{dt} = -k_{\max} Q$$

$$\log\left(\frac{N}{N_0}\right) = \log(\exp(-k_{\max} t)) \frac{1 + Q(0)}{1 + Q(0) \exp(-k_{\max} t)}$$

Q – variável relacionada com o estado fisiológico dos microrganismos

Modelos de inativação

❖ primários



Gompertz

Bhaduri et al (1991)
Linton et al. (1995, 1996)
Xiong et al. (1999)



Listeria monocytogenes


$$\log N = \log N_0 - \log \left(\frac{N_0}{N_{res}} \right) \exp \left(- \exp \left(\frac{k e}{\log \left(\frac{N_0}{N_{res}} \right)} (L - t) + 1 \right) \right)$$



reparameterização para inativação com base em Zwietering (1990)

Modelos de inativação

❖ primários



Gompertz

Bhaduri et al (1991)
Linton et al. (1995, 1996)
Xiong et al. (1999)



Listeria monocytogenes

$$\log N = \log N_0 - \log \left(\frac{N_0}{N_{res}} \right) \exp \left(- \exp \left(\frac{k e}{\log \left(\frac{N_0}{N_{res}} \right)} (L - t) + 1 \right) \right)$$



reparameterização para inativação com base em Zwietering et al. (1990)

Logística

$$\log N = \frac{c}{1 + \exp(k(t - L))}$$

c – constante


Exemplos

Data of *L. monocytogenes* Scott A at 52,56,60,64,68°C

(24 hours incubation at 5°C in half cream)

Casadei et al. (1998)

Gompertz



Gil (2002)


Statistica 6.0

Exemplos

Data of *L.monocytogenes* Scott A at 52,56,60,64,68°C
(24 hours incubation at 5°C in half cream)

Casadei et al. (1998)

Gompertz



Gil (2002)

Statistica 6.0


Exemplos

Data of *L. monocytogenes* Scott A at 52,56,60,64,68°C

(24 hours incubation at 5°C in half cream)

Casadei et al. (1998)

Logística




Exemplos

Data of *L.monocytogenes* Scott A at 52,56,60,64,68°C
(24 hours incubation at 5°C in half cream)

Casadei et al. (1998)

Logística




Gil (2002)

Statistica 6.0

Modelos matemáticos

❖ primários



❖ secundários
parâmetros

Modelos matemáticos

❖ secundários

Arrhenius

$$k = k_0 \exp\left(-\frac{E_a}{RT}\right) \rightarrow \ln k = \ln k_0 - \frac{E_a}{RT}$$

$$k = k_{\text{ref}} \exp\left(-\frac{E_a}{R} \left(\frac{1}{T} - \frac{1}{T_{\text{ref}}}\right)\right)$$

Davey / Arrhenius modificado

$$\ln k = C_0 + \frac{C_1}{T} + \frac{C_2}{T^2} + C_3 a_w + C_4 a_w^2$$

“Square-root type models”

Ratkowsky *et al.* (1982)

$$\sqrt{k} = b(T - T_{\text{min}})$$

McMeekin *et al.* (1987)

$$\sqrt{k} = b(T - T_{\text{min}}) \sqrt{(a_w - a_{w\text{min}})}$$

Adams *et al.* (1991)

$$\sqrt{k} = b(T - T_{\text{min}}) \sqrt{(\text{pH} - \text{pH}_{\text{min}})}$$

McMeekin *et al.* (1992)


$$\sqrt{k} = b(T - T_{\text{min}}) \sqrt{(a_w - a_{w\text{min}})} \sqrt{(\text{pH} - \text{pH}_{\text{min}})}$$

min – valor mínimo para crescimento

Exemplos

Data of *L.monocytogenes* Scott A
(24 hours incubation at 5°C in half cream)

Arrhenius



$k = f(T)$

Gompertz

$$k=0.0216 \exp(-203.3/R^*(1/T-1/333.15))$$

DEa=28.85 kJ/mol

DK_{ref}=4.58x10⁻³ s⁻¹

Logística

$$k=0.0337 \exp(-206.6/R^*(1/T-1/333.15))$$

DEa=27.56 kJ/mol

DK_{ref}=7.31x10⁻³ s⁻¹

Gil (2002)


Statistica 6.0

Exemplos

Data of *L.monocytogenes* Scott A
(24 hours incubation at 5°C in half cream)

Gompertz

atraso = $f(T)$



$DE_a = 7.485 \text{ kJ/mol}$

$DL_{ref} = 7.595 \text{ s}^{-1}$

Gil (2002)


Statistica 6.0

Exemplos

Data of *L.monocytogenes* Scott A
(24 hours incubation at 5°C in half cream)

Logística

atraso = $f(T)$



$DE_a = 3.591 \text{ kJ/mol}$

$DI_{ref} = 6.154 \text{ s}^{-1}$

Gil (2002)

Statistica 6.0

Modelos matemáticos

Condições variáveis de

temperatura

pH

a_w



maior complexidade !!



$$\frac{d(\log N)}{d(\text{tempo})}$$

Modelos matemáticos

Gompertz

situação dinâmica de temperatura

$$\downarrow \frac{d(\log N)}{d(\text{tempo})}$$

$$\log N = \log N_0 - \int_0^t \left[k \exp(1) \exp \left(\frac{k \exp(1)}{\log \left(\frac{N_0}{N_{\text{res}}} \right)} (L - t') + 1 \right) \exp \left(- \exp \left(\frac{k \exp(1)}{\log \left(\frac{N_0}{N_{\text{res}}} \right)} (L - t') + 1 \right) \right) \right] dt'$$

$$k = k_{\text{ref}} \exp \left(- \frac{E_a}{R} \left(\frac{1}{T} - \frac{1}{T_{\text{ref}}} \right) \right)$$

$$L = a \exp \left(b \left(\frac{1}{T} - \frac{1}{T_{\text{ref}}} \right) \right)$$

Modelos de inactivação

❖ terciários

softwares

crescimento de microrganismos
previsão de tempo de prateleira

inactivação de microrganismos

lacunas

Limitações

- interacções entre microrganismos
- diversidade natural das estirpes
- estrutura complexa dos alimentos
- alimento/microrganismo
- modelagem da fase 'atraso'
- modelagem da fase 'cauda'
- previsões reais com condições ambientais variáveis

Envolvimento da equipa num projecto internacional ...

2001-2004

BUGDEATH

www.frperc.bris.ac.uk/bugdeath.htm

microbiologia
predictiva

pasteurização de
alimentos à
superfície

fronteira importante de contaminação

"BUGDEATH" (QLRT-2001-01415)



Micrograph of a Salmonella bacteria

Predicting the reduction in microbes on the surface of foods during pasteurisation.

- [Tell me more about the Bugdeath project.](#)
- [What are the objectives of the Bugdeath project?](#)
- [Who are the project partners?](#)
- [Meetings](#)
- [Industrial Advisory Group \(IAG\)](#)
- [Partner area of web site \(specifically for project partners - password needed to enter\)](#)

Bugdeath

BUGDEATH is a research project funded by the European Commission to produce accurate predictive models of the reductions in microbial numbers that can be achieved on the surface of foods during surface pasteurisation processes.

Food poisoning is increasing throughout the European Union (EU). Over 60% of outbreaks are associated with meat, fresh fruit and salad vegetables. Most of the contamination by pathogenic and spoilage organisms is present on the surface of foods at the time of harvesting or is transferred to the surfaces during slaughter and processing. Accurate microbial death models would be of considerable help to the food industry in the development of surface pasteurisation systems for meat, fruit and vegetables. This will in turn lead to safer foods with improved quality and shelf life.



Personnel involved in Bugdeath from left to right

Bugdeath

Parceiros

1. Food Refrigeration and Process Engineering Research Center (**FRPERC**)
University of Bristol
2. Escola Superior de Biotecnologia (**ESB**)
Universidade Católica Portuguesa
3. Department of Chemical Engineering
Katholieke Universiteit Leuven (**KUL**)
4. Teagasc, The National Food Center (**NFC**)
5. Campden & Chorleywood Food Research Association (**CCFRA**)
6. Faculty of Applied Science
University of West England (**UWE**)
7. Laboratoire de Génie des Procédés Alimentaires (**ENITIA**)
Ecole Nationale d'Ingénieurs des Techniques des Industries Agricoles et Alimentaires
8. Institute National de la Recherche Agronomique (**INRA**)

Bugdeath

Objectivos

- estudos de inactivação à superfície de alimentos sólidos
- desenvolvimento de modelos de inactivação precisos e exactos
- desenvolvimento de equipamento para pasteurização à superfície
'rig apparatus'
- desenvolvimento de software para previsão

tarefa da ESB

Bugdeath

alimentos estudados

batata

frango

bife

microrganismos

Listeria monocytogenes

E. coli

Salmonella

Campylobacter

rig apparatus



Bugdeath


Bugdeath

processo térmico

- vapor
- ar seco

história de temperatura

rig apparatus



podem ser
testadas
diferentes
histórias de
temperatura

Escola Superior de Biotecnologia, Universidade Católica Portuguesa

Team

Pedro Pereira
Maria Manuel Gil
Teresa Brandão
Cristina Silva


Sessão de disseminação
Chipping Campden, Agosto 2004

Development of a user-friendly
combined heat, mass transfer and
microbial death model




Objectives

- Create a software application



User-friendly tool




foods

Simulation of inactivation kinetics of
microorganisms on the surface of
during pasteurisation treatments



Introduction

- Need to obtain reliable data on relationship between microbial death and the surface temperature of real foods




Lead to the design,
construction and
commission of equipment

Software application to
simulate results obtained in the
rig apparatus



Modelling



Modelling

two modelling approaches

- Accurate modelling of heat transfer
 - Description of the phenomena induced to the food surface by a thermal process
- Modelling microbial inactivation behaviour under such temperature conditions



Heat transfer model

estimation of surface temperature

- One dimensional heat transfer model
- Combination of different phenomena
 - Conduction
 - Convection
 - Evaporation/condensation of water or steam
 - Radiation (not considered)



Heat transfer model

- Simplified model

Conduction/convection - Fourier

Geometry – plane sheet

No mass transfer phenomena considered




- Limits

Air velocity ranged from 15 m/s to 25 m/s

Extrapolation locked outside of conditions that can be verified in the test rig




Heat transfer model



Boundary conditions - bottom

Exchanges between the bottom of the product and the support can be neglected – product thickness < 0.5 cm



Experimentally observed in the rig



Heat transfer model

$$\lambda \left(\frac{\partial T}{\partial x} \right)_{\text{top}} = h_{\text{eff}} (T_{\text{air or steam}} - T_{\text{sur}})$$

(3)


(1)

$$\frac{\partial T(x, t)}{\partial t} = \alpha \frac{\partial^2 T(x, t)}{\partial x^2}$$

Crank-Nicholson method

$$\lambda \left(\frac{\partial T}{\partial x} \right)_{\text{bot}} = h_{\text{inf}} (T_{\text{sup}} - T_{\text{inf}})$$

(2)



- t – time
- T – temperature
- h – heat transfer coefficient
- α – thermal diffusivity
- λ – Thermal conductivity



Effective coefficient – dry conditions

$$h_{\text{eff}} = h \frac{T_{\text{air}} - T_{\text{sur}}}{T_{\text{max}} - T_{\text{sur}}} + K_m \Delta H \frac{P_T - a_{w_{\text{sur}}} P_{T_{\text{sur}}}}{T_{\text{max}} - T_{\text{sur}}}$$


convection evaporation

k_m – mass transfer coefficient

ΔH – latent heat of water evaporation



Effective coefficient – wet conditions (steam)



Due to complexity of mathematical expressions to be incorporated in the software

$$T_{\text{sur}} = T_{\text{steam}} - 3^{\circ}\text{C}$$



Experimentally observed in the rig



Microbial inactivation model

- To predict microbial content at the surface of foods
- It has the advantage of dealing with time-varying temperature conditions
- Typical of pasteurisation treatments



Microbial inactivation model

$$\frac{dN}{dt} = -k_{\max} k_Q(Q)N$$


$$\frac{dQ}{dt} = -k_{\max} Q$$

N – microbial cell density

Q – variable related to the physiological state of the cells

k_{\max} – inactivation rate constant

$$\log \frac{N}{N_0} = \log(\exp(-k_{\max} t)) \frac{1 + Q(0)}{1 + Q(0) \exp(-k_{\max} t)}$$



Microbial inactivation model

$$k_{\max}(T, a_w) = \frac{\ln 10}{D_{ref}} \exp\left(\frac{\ln 10}{z} (T - T_{ref})\right) \exp\left(\frac{\ln 10}{z_{a_w}} (a_w - 1)\right) + c1$$



$T = T_{sur} \rightarrow$ Calculated on the basis of all considerations of heat transport



Software Program



'Bugdeath' - funded by the European Commission under the EC Framework 5;
Quality of Life and Management of Living Resources Programme



Software Program


- The programme was developed using Real Basic® 5.2 application
- Food/microorganism selection is allowed
(database of thermal properties and kinetic parameters)
- On the basis of the selection of a heating regime of the medium, the programme allows prediction of the food surface temperature and simulates the microbial load content along the whole process time




First screen – product/microorganism

The screenshot shows the 'BugDeath' software interface. The window title is 'BugDeath'. There are three tabs: 'Product', 'Process', and 'Output'. The 'Product' tab is active. Below the tabs, there are two main sections: 'Product' and 'Microorganism'. The 'Product' section includes a dropdown menu for 'Product' (set to 'Beef'), input fields for 'Diffusivity' (1.23e-07), 'Conductivity' (4.5e-01), 'Product thickness (m)' (0.016), 'Init. Product Temp. (°C)' (13.4), 'Sample Diameter (m)' (0.05), and 'Water activity' (0.45). The 'Microorganism' section includes a dropdown menu for 'Microorganism' (set to 'S. typhimurium'), input fields for 'D (min)' (6.6e+1), 'Safe Count' (1.e+0), 'Initial Count' (10000000), 'Z (°C)' (7), 'Z(aw)' (0.2), 'R. Temp. (°C)' (57.2), 'Z(aw)' (0.2206), and 'R. Temp. (°C)' (2.279372077). There are also labels 'C1' and 'Q0' at the bottom of the input fields. On the right side, there are three tabs: 'Temperature variation', 'Microbial load variation', and 'Data Table'. The 'Temperature variation' tab is active, showing a large empty area. At the bottom right of the window, there are labels 'Surface Temperature', 'Medium Temperature', and 'Temperature'.


'Bugdeath' - funded by the European Commission under the EC Framework 5;
Quality of Life and Management of Living Resources Programme




Second screen – process




'Bugdeath' - funded by the European Commission under the EC Framework 5;
Quality of Life and Management of Living Resources Programme




Output – graphic/temperature




'Bugdeath' - funded by the European Commission under the EC Framework 5;
Quality of Life and Management of Living Resources Programme



Output – graphic/microbial load



'Bugdeath' - funded by the European Commission under the EC Framework 5;
Quality of Life and Management of Living Resources Programme



Output – data table

The screenshot shows the 'BugDeath' software interface. The 'Data Table' tab is active, displaying the following data:

Time (s)	Medium Temp. (°C)	Surface Temp. (°C)	Microbial Load
0	0	14.02522	7
0.6081229	100	39.29868	6.991117
1	100	31.74754	6.991117
1.608123	100	35.68345	6.991117
2	100	36.00534	6.991117
2.608123	100	36.9512	6.991117
3	100	38.14414	6.991117
3.608123	100	38.62585	6.991117
4	100	39.78456	6.991117
4.608123	100	40.16022	6.991117
5	100	41.19636	6.991117
5.608123	100	41.524	6.991117
6	100	42.4504	6.991117
6.608123	100	42.74452	6.991117
7	100	43.58093	6.991117
7.608123	100	43.84836	6.991117
8	100	44.61078	6.991117
8.608123	100	44.85613	6.991117
9	100	45.55682	6.991117
9.608123	100	45.78353	6.991117
10	100	46.43196	6.991117
10.60812	100	46.64271	6.991116
11	100	47.2463	6.991116
11.60812	100	47.44323	6.991116
12	100	48.0079	6.991116
12.60812	100	48.19275	6.991116
13	100	48.72329	6.991115
13.60812	100	48.89746	6.991115
14	100	49.39783	6.991115

'Bugdeath' - funded by the European Commission under the EC Framework 5;
Quality of Life and Management of Living Resources Programme



Conclusions / outputs

- Software application simulates the results obtained in the rig apparatus
- Valuable for developing appropriate and safety thermal processes
- Marketed and commercially available
- Educational purposes (simulation of real food processes)



Workshop

Predictive Microbiology

Aplicações e Novas Fronteiras

Obrigada

Teresa Brandão

Cristina Silva



UNIVERSIDADE CATÓLICA PORTUGUESA
Escola Superior de Biotecnologia

4 de Março 2005