



SOCIEDADE
PORTUGUESA
DE QUÍMICA

ANALÍTICA 2018


9th Meeting of Division of Analytical Chemistry

26-27 MARCH, PORTO-PORTUGAL

FFUP/ICBAS – UNIVERSITY OF PORTO

www.analitica2018.eventos.chemistry.pt

**BOOK OF
ABSTRACTS**



PROTEIN-PHENOLIC INTERACTION AS A STRATEGY TO REDUCE THE PRECURSORS OF VOLATILE PHENOLS IN WINE

Susana S. M. P. Vidigal, Francisco M. Campos, José A. Couto,
António O. S. S. Rangel, Tim Hogg

*Universidade Católica Portuguesa, CBQF - Centro de Biotecnologia e Química Fina –
Laboratório Associado, Escola Superior de Biotecnologia, Rua Arquiteto Lobão Vital 172,
4200-374 Porto, Portugal. svidigal@porto.ucp.pt*

In the present work, the interaction between phenolic acids (caffeic acid, *p*-coumaric acid and ferulic acid) and proteins was studied as a potential strategy to reduce the content of the precursors of volatile phenols in wine. These phenolics result from the contamination of the wine by *Brettanomyces bruxellensis* causing a wine defect often referred as the “*Brett character*” [1]. In this context, the aim of the present work is the development of a biological strategy, in which the precursors can be removed from the wine or otherwise be rendered unreactive and for that reason be unavailable for conversion into volatile phenols.

Due to their possible beneficial effects on human health, the interaction of proteins with plant-derived phenolic compounds has been an object of study [2]. This interaction can be studied using a fluorescence quenching approach [3], which has been widely employed to characterise the biochemical interactions between these types of molecules.

Bovine serum albumin (BSA) was used as a model protein and the fluorescence of the protein was measured in the absence and in the presence of the quencher - the phenolic acid. This protein-phenolic interaction can be affected by some environmental conditions, such as medium composition, pH, temperature, and protein concentration [2, 4]. The effect of the environmental conditions on the interaction between BSA and caffeic, coumaric and ferulic acids was evaluated by the determination of the binding parameters, K_a and n , where K_a is the binding constant and n is the number of binding sites per phenol/protein. Higher values were obtained for the interaction of caffeic acid with BSA.

Acknowledgements: Susana Vidigal acknowledges the financial support in the framework of the project “Biological tools for adding and defending value in key agro-food chains (bio – n2 – value)”, nº NORTE-01-0145-FEDER-000030, funded by Fundo Europeu de Desenvolvimento Regional (FEDER), under Programa Operacional Regional do Norte - Norte2020. The authors would also like to thank the scientific collaboration of the project UID/Multi/50016/2013.

- [1] Suárez, R.; Suárez-Lepe, J. A.; Morata, A.; Calderón, F. *Food Chem.* **2007**, *102*, 10-21.
- [2] Ozdal, T.; Capanoglu, E.; Altay, F. *Food Res. Int.* **2013**, *51*, 954-970.
- [3] Le Bourvellec, C.; Renard, C. M. G. C. *Crit. Rev. Food Sci. Nutr.* **2012**, *52*, 213-248.
- [4] Czubinski, J.; Dwiecki, K. *Int. J. Food Sci. Tech.* **2017**, *52*, 573-585.

TÍTULO

Book of Abstracts Analítica 2018 - 9th Meeting
of Division of Analytical Chemistry

EDITORIAL BOARD

Marcela A. Segundo
Eduarda M. P. Silva
Luísa Barreiros
Lúcia Saraiva

TYPESET AND GRAPHIC DESIGN

Joana Macedo, FFUP

ALL RIGHTS RESERVED

This publication may not be reproduced or
transmitted in whole or in part by any process,
electronic, mechanical, photocopying, recording
or otherwise, without the prior permission of the
editorial board.

EDITION

Sociedade Portuguesa de Química
Av. da República, 45 - 3^o Esq
1050-187 Lisboa - Portugal
ISBN: 978-989-8124-21-0

DATE

March 2018