

Effect of Ultrasonication, Thermo-sonication and UV-C Irradiation on the Quality of Strawberries (*Fragaria ananassa*) and Red Bell Peppers (*Capsicum annuum* L.)



D. M. SANTOS PEDRO, E.M.C. ALEXANDRE, J. FUNDO, T.R.S. BRANDÃO, C. L.M. SILVA (*)

Escola Superior de Biotecnologia - Universidade Católica Portuguesa
Rua Dr. António Bernardino de Almeida, 4200-072 Porto, Portugal; Fax: +351 225 090 351, www.esb.ucp.pt/lopa

(*) E-mail: dsilva@esb.ucp.pt



Introduction

Alternative non-thermal technologies constitute emergent challenging methods aiming at reducing pernicious effects of thermal methods, by preserving quality and nutritional attributes of fruits and vegetables, and yielding safe and less-perishable products. The application of ultrasounds and ultraviolet (UV-C) irradiation are examples of non-thermal technologies that may have potential applications in the food industry.

Ultraviolet (UV) light occupies a wide band of wavelengths in the non-ionising region of the electromagnetic spectrum between X-rays (200 nm) and visible light (400 nm) (Hollósy, 2002). The germicidal range is in the region of short-wave UV (UV-C), with wavelengths between 200 and 280 nm, being the 254 nm the most lethal (Summerfelt, 2003). Exposure to low doses of UV-C has been reported to reduce post-harvest decay of fruits and vegetables (Lu *et al.*, 1987; Erkan *et al.*, 2001; Marquerie *et al.*, 2002; Allende and Artés, 2003), which accredits prospective application of UV-C in the postharvest industry.

Ultrasound is defined as pressure waves with a frequency of 20 kHz or more (Butz and Tauscher, 2002). Ultrasound may be used at frequencies between 20 kHz and 10 MHz. Higher-power ultrasound at lower frequencies (20-100 kHz) has the ability to cause cavitation, which has the capacity to inactivate microbes and enzymes (Knorr *et al.*, 2002, 2004; Piyasena *et al.*, 2003). The conjoint application of mild temperatures may enhance the ultrasonication effect (thermosonication), especially in terms of product's safety (Mason *et al.*, 1996; López-Malo *et al.*, 2005), with minor changes in terms of quality parameters, when compared to conventional thermal methods.

Objective

The objective of this work was to study the effect of ultrasonication, thermo-sonication and UV-C irradiation on the quality attributes of strawberries (*Fragaria ananassa*) and red bell peppers (*Capsicum annuum*, L.).

Materials and Methods

Strawberries (*Fragaria ananassa*) and red bell peppers (*Capsicum annuum*, L.) were acquired in a local market. They were pre-washed and carefully soaked with an absorbant towel. Strawberries were used either as whole, for ultrasonication and thermo-sonication studies, or cut into two equal halves, for the UV-C irradiation treatments or texture analysis studies. Red bell peppers were cut into pieces of 1x2 cm for all treatments.

UV-C treatments were carried out in an UV-C chamber (Figure 1) that was designed by Universidade do Algarve (EST-Ualg, Faro, Portugal). Samples were carefully placed in petri plates and submitted to a bank of four germicidal UV lamps (TUV G30T8, 16 W, Philips, peak emission at 254 nm) for 2 minutes, at an average intensity of 12,36W/m². The intensity of flux of lamps and dose of exposure (*time x intensity*) were continuously measured using an UV digital photometer (DO 9721 Delta Ohm).

Ultrasonication and thermo-sonication treatments were carried out in an ultrasound machine (Figure 1) (Bandelin Sonorex RK 100H) operating at 32 kHz. Samples were treated for 2 minutes at 20°C (ultrasonication), and 50°C and 65°C (thermosonication). Control water treatments at the same temperatures were also performed.




Figure 1 – Equipments used for (A.1-2) UV-C irradiation and (B.1-2) ultra- and thermo-sonication treatments.

The quality factors analysed were colour and texture for both products, in addition to anthocyanins content in the case of strawberries. At least 4 replicates of at least 30 samples each were analyzed.

Colour was measured using the Hunter (L, a and b) scale, with a colorimeter (CR-300, Minolta). The total colour difference (TCD) = $[(L-L_0)^2 + (a-a_0)^2 + (b-b_0)^2]^{1/2}$; being index "0" indicative of initial reference values of fresh product) was the parameter considered for evaluation of colour.

Anthocyanins were assayed by a spectrophotometric procedure (spectrophotometer UV-1601, Shimadzu). Results were obtained in terms of monomeric pigment anthocyanin (pelargonidin 3-glycoside).

Texture analysis was conducted in a texturometer (TA-XT2plus, TA Instruments) equipped with a 5 kg force cell. Compression mode tests (30% deformation, 1mm/s velocity) were performed using a 10 mm or a 36 mm diameter cylindrical probe for red bell peppers or strawberries, respectively. Hardness values were registered as the maximum force observed during compression of samples.

The experimental results were compared by analysis of variance (two-way ANOVA, Analysis Tool Package, Excel 2000, Microsoft®, USA).

Results and Discussion

Results of strawberries and red bell peppers quality are, respectively, presented in tables 1 and 2.

For both products, irradiation with UV-C light constituted the most harmless treatment, with little changes on colour and texture (hardness).

After ultrasonication (US 20°C) samples showed distinct colour in relation to fresh. However, this effect was very small when compared to the effect of thermo-sonication, especially at 65°C. At this temperature, thermo-sonication caused great differences on the colour of samples. Water treatments carried out at the same temperature showed very similar results, which suggests that the effect on colour is due to the temperature effect, and not exclusively to the sonication of samples.

Table 1. Results for colour, hardness and anthocyanins of strawberries (*Fragaria ananassa*) treated by ultraviolet radiation (UV-C), ultrasonication (US 20°C), thermo-sonication (US 50°C and US 65°C) and water (W 20°C, W 50°C and W 65°C).

	Color (TCD ^a)	Hardness (normalized values ^b)	Anthocyanins (normalized values ^b)
UV-C	1,562 ± 0,829	1,025 ± 0,206	0,888 ± 0,076
US 20°C	2,626 ± 0,235	0,999 ± 0,070	1,082 ± 0,083
W 20°C	2,864 ± 0,355	0,763 ± 0,129	1,250 ± 0,010
US 50°C	4,620 ± 0,463	0,919 ± 0,246	0,866 ± 0,180
W 50°C	3,084 ± 0,243	0,827 ± 0,154	0,945 ± 0,080
US 65°C	6,387 ± 0,610	0,640 ± 0,233	0,835 ± 0,351
W 65°C	4,347 ± 0,416	0,279 ± 0,079	0,756 ± 0,112

^a TCD – Total Color Difference

^b normalized values in relation to untreated samples

Table 2. Results for colour and hardness of red bell peppers (*Capsicum annuum*) treated by ultraviolet radiation (UV-C), ultrasonication (US 20°C), thermo-sonication (US 50°C and US 65°C) and water (W 20°C, W 50°C and W 65°C).

	Color (TCD ^a)	Hardness (normalized values ^b)
UV-C	1,344 ± 0,853	0,971 ± 0,306
US 20°C	4,778 ± 1,307	1,146 ± 0,451
W 20°C	3,051 ± 1,034	1,070 ± 0,414
US 50°C	5,528 ± 1,856	1,214 ± 0,500
W 50°C	3,144 ± 1,000	0,910 ± 0,420
US 65°C	6,572 ± 1,770	1,076 ± 0,480
W 65°C	7,487 ± 1,525	0,587 ± 0,282

^a TCD – Total Color Difference

^b normalized values in relation to untreated samples

In the case of strawberries a slight decrease in anthocyanins content was observed at higher temperatures, which may be related with observed total colour differences.

Interesting results were obtained in relation to the products hardness. In the case of strawberries (table 1), samples turned softer with increasing of temperature. However, the decrease of hardness was much more pronounced on the water treated. In the case of red bell peppers (table 2), softening of samples was only observed for water treated samples at 65°C. These results suggest that, somehow sonication of samples (at the used frequency), even when coupled with temperature, seems to preserve the structure of vegetable tissues, preventing them from texture loss when higher temperatures are used.

Conclusions

- UV-C constituted a harmless treatment, with little or no effect on colour, texture or anthocyanins content.
- Ultrasonication caused small colour changes and no significant effects on textures of both products.
- Thermo-sonication at 65°C caused significant changes on colour of strawberries and red bell peppers.
- When compared to water treated samples, thermo-sonicated ones showed higher texture retention. This effect was much more pronounced on red bell peppers.

References

- Allende A. and Artés, F. (2003). UV-C radiation as a novel technique for keeping quality of fresh processed Lollo Rosso lettuce. *Food Research International*, 36(8):739-756
- Butz, P. and Tauscher, B. (2002). Emerging technologies: Chemical aspects. *Food Research International* 35(2-3):279-284
- Erkan, M., Wang, C.Y. and Krizek, D.T. (2001). UV-C radiation reduces microbial populations and deterioration in *Cucurbita pepo* fruit tissues. *Environmental and Experimental Botany*, 45(1):1-9
- Hollósy, F. (2002). Effects of ultraviolet radiation on plant cells. *Micron*, 33(2):179-197
- Knorr, D., Ade-Omowaye, B.J. and Heinz, V. (2002). Nutritional improvement of plant foods by non-thermal processing. *Proceedings of the Nutrition Society*, 61(2):311-318
- Knorr, D., Zenker, M., Heinz, V. and Lee, D.-U. (2004). Applications and potentials of ultrasonics in food processing. *Trends in Food Science and Technology*, 15(5):261-266
- López-Malo, A., Palou, E., Jiménez-Fernández, M., Altamora, S.M., Guerrero, S. (2005). Multifactorial fungal inactivation combining thermo-sonication and antimicrobials. *J. Food Engineering*, 67(1-2):87-93
- Lu, J.Y., Stevens, C., Yakubu, P., Loretan, P.A. and Eskin, D. (1987). Gamma, electron beam and UV radiation on control of storage roots and quality of Walla Walla onions. *J. Food Processing and Preservation*, 12:53-62
- Marquerie, D., Michiels, C.W., Geeraerd, A.H., Schenk, A., Soontjen, C., Van Impe, J.F. and Nicolai, B.M. (2002). Using survival analysis to investigate the effect of UV-C and heat treatment on storage rot of strawberry and sweet cherry. *International Journal of Food Microbiology*, 73(2-3):187-196
- Mason, T.J., Paniwnyk, L. and Lorimer, J.P. (1996). The uses of ultrasound in food technology. *Ultrasounds and sonochemistry*, 3(3):S253-S260
- Piyasena, P., Mohareb, E. and McKellar, R.C. (2003). Inactivation of microbes using ultrasound: a review. *International Journal of Food Microbiology* 87(3):207-216
- Summerfelt, S.T. (2003). Ozonation and UV irradiation – an introduction and examples of current applications. *Aquacultural Engineering*, 28(1-2):21-36

Acknowledgments

The authors acknowledge the financial support through Programa Operacional Agricultura e Desenvolvimento Rural – Projecto AGRO nº822 (Novas Tecnologias de Processamento de Hortofrutícolas Congeladas – EMERCON). The authors Alexandre E. M. C., Santos Pedro D. M. and Brandão T. R. S. would like to thank Fundação para a Ciência e a Tecnologia (grants SFRH/BD/16042/2004, SFRH/BPD/9174/2002 and SFRH/BPD/11580/2002, respectively).